

SUPERCOHERENT STATES, SUPER KÄHLER GEOMETRY AND GEOMETRIC QUANTIZATION

Amine M. El Gradechi^{1,2,*}, Luis M. Nieto^{1,†}

¹⁾ *Centre de Recherches Mathématiques, Université de Montréal
C.P. 6128-centre-ville, Montréal (Québec) H3C 3J7, Canada*

²⁾ *Department of Mathematics and Statistics, Concordia University
Montréal (Québec) H4B 1R6, Canada*

ABSTRACT

Generalized coherent states provide a means of connecting square integrable representations of a semi-simple Lie group with the symplectic geometry of some of its homogeneous spaces. In the first part of the present work this point of view is extended to the supersymmetric context, through the study of the $\mathrm{OSp}(2/2)$ coherent states. These are explicitly constructed starting from the known abstract typical and atypical representations of $\mathrm{osp}(2/2)$. Their underlying geometries turn out to be those of supersymplectic $\mathrm{OSp}(2/2)$ -homogeneous spaces. Moment maps identifying the latter with coadjoint orbits of $\mathrm{OSp}(2/2)$ are exhibited via Berezin's symbols. When considered within Rothstein's general paradigm, these results lead to a natural general definition of a super Kähler supermanifold, the supergeometry of which is determined in terms of the usual geometry of holomorphic Hermitian vector bundles over Kähler manifolds. In particular, the supergeometry of the above orbits is interpreted in terms of the geometry of Einstein-Hermitian vector bundles. In the second part, an extension of the full geometric quantization procedure is applied to the same coadjoint orbits. Thanks to the super Kähler character of the latter, this procedure leads to explicit super unitary irreducible representations of $\mathrm{OSp}(2/2)$ in super Hilbert spaces of L^2 superholomorphic sections of prequantum bundles of the Kostant type. This work lays the foundations of a program aimed at classifying Lie supergroups' coadjoint orbits and their associated irreducible representations, ultimately leading to harmonic superanalysis. For this purpose a set of consistent conventions is exhibited.

Key-Words: Coherent states, supersymplectic geometry, super Kähler supermanifolds, geometric quantization, Lie superalgebras

1991 MSC: 17A70, 32C11, 53C55, 58A50, 58F06, 81R30, 81S10

PACS: 02.20 Sv, 02.40, 03.65 Sq

*E-mail: elgrade@ere.umontreal.ca

†E-mail: nietol@ere.umontreal.ca

RÉSUMÉ

Les états cohérents généralisés fournissent un moyen de relier les représentations de carré intégrables d'un groupe de Lie semi-simple à la géométrie symplectique de certains de ses espaces homogènes. Dans la première partie de ce travail, l'extension supersymétrique de ce point de vue est considérée à travers l'étude des états cohérents de $\mathrm{OSp}(2/2)$. Ceux-ci sont explicitement construits à partir des représentations abstraites typique et atypique connues de $\mathrm{osp}(2/2)$. Les géométries associées à ces états se révèlent être celles d'espaces homogènes supersymplectiques de $\mathrm{OSp}(2/2)$. Des applications moments identifiant ces derniers avec des orbites coadjointes de $\mathrm{OSp}(2/2)$ sont produites via les symboles de Berezin. Interprétés dans le cadre de la théorie générale développée par Rothstein, ces résultats conduisent à une définition générale et naturelle d'une supervariété super Kählerienne. La supergéométrie de celle-ci est déterminée en termes de la géométrie habituelle de fibrés vectoriels holomorphes Hermitiens au dessus de variétés Kähleriennes. En particulier, la supergéométrie des orbites mentionnées ci-dessus s'interprètent en termes de fibrés d'Einstein-Hermitiens. Dans la seconde partie, une extension de la procédure complète de quantification géométrique est appliquée aux mêmes orbites coadjointes. Leur caractère super Kählerien permet alors de produire des représentations super unitaires irréductibles explicites de $\mathrm{OSp}(2/2)$ dans des super espaces de Hilbert de sections L^2 superholomorphes de fibrés préquantiques du type décrit par Kostant. Ce travail pose les premiers jalons d'un programme ayant pour objectif de classifier les orbites coadjointes des supergroupes de Lie et les représentations qui leurs sont associées, pour finalement aboutir à une superanalyse harmonique. Pour cela un ensemble consistant de conventions est présenté.

Contents

1	Introduction	1
2	$\mathfrak{osp}(2/2)$ representations and typical $\mathrm{OSp}(2/2)$-CS	5
2.1	$\mathfrak{osp}(2/2)$ representations	5
2.2	$\mathrm{OSp}(2/2)$ coherent states	12
3	Supersymplectic geometry and the $\mathrm{OSp}(2/2)$-CS	14
3.1	The supersymplectic form	15
3.2	The classical observables	16
3.3	The Liouville supermeasure	19
4	More about supergeometry	20
4.1	Supersymplectic supermanifolds	21
4.2	$\mathcal{D}^{(1 2)}$ as a holomorphic supermanifold	23
4.3	$\mathcal{D}^{(1 2)}$ as a super Kähler supermanifold	24
5	Geometric quantization	27
5.1	Super prequantization	29
5.2	Super polarization	30
6	Atypical $\mathrm{OSp}(2/2)$-CS and associated orbits	35
7	Miscellaneous results and discussions	38
8	Conclusions and outlook	41
A	Conventions and notation	42
B	$\mathfrak{osp}(2/2)$ representations: more details	44
	References	48

1 Introduction

1.1. Coherent states were originally those very special quantum states of the harmonic oscillator first introduced by Schrödinger [1] and studied further by Glauber [2]. Their special character stems from their property of being the closest possible states to the classical theory. This is reflected in the fact that they minimize Heisenberg's uncertainty relations. Moreover, they form an overcomplete basis of the Hilbert space of quantum states. Since Glauber's contribution, the concept of coherent states has evolved very rapidly. The key step of this evolution was without any doubt Perelemov's group theoretical generalization [3]. Nowadays' coherent states find applications in several areas of physics and mathematics. Let us mention, for instance, their occurrence in quantum optics [4], in signal analysis (where they are called wavelets) [5], and in mathematical physics (in connection with the quantization-versus-classical-limit procedures) [6, 7] (see also [3] and references therein). This last application constitutes the main interest of the present paper. More precisely, we extend to the supersymmetric context both the coherent states approach to the evaluation of the classical limit, and the geometric quantization of coadjoint orbits. Even though these two points have already been studied in the case of the $\mathrm{OSp}(1/2)$ Lie supergroup in [8, 9], the analysis of a less trivial example, such as the $\mathrm{OSp}(2/2)$ Lie supergroup considered here, brings new insights that improve our understanding of the very interesting supergeometric structures underlying the so-called supercoherent states. Let us now present an overview of the subject. More details and references about the coherent states methods can be found in [3, 4, 10].

1.2. The harmonic oscillator coherent states admit a group theoretical construction based on the Weyl-Heisenberg group underlying this physical system. By extending this construction to general Lie groups, Perelomov introduced the notion of generalized coherent states [3]. For a given Lie group G and a given unitary irreducible representation U of G in some Hilbert space $\mathcal{H}$, the generalized coherent states are the states belonging to a $U(G)$ -orbit in $\mathcal{H}$ through a chosen initial state, usually called the fiducial state. The generalized coherent states so defined are too general to share the interesting properties of the harmonic oscillator ones. Indeed, the overcompleteness property holds only when U is a square integrable representation, and the uncertainty relations associated with the Lie algebra $\mathfrak{g}$ of G are minimized only when the fiducial state is a highest weight state [3, 11, 12]. In what follows we will only consider this type of generalized coherent states and we will simply designate them by coherent states (CS) or G -coherent states (G -CS).

1.3. Not all Lie groups possess square integrable representations. Unfortunately, this is the case for some physically relevant groups, such as the Poincaré group. There is however enough room to actually take advantage of the particularly rich properties of the generalized CS. Indeed, all representations of compact semi-simple Lie groups, and all discrete series representations of the non-compact semi-simple Lie groups, are square integrable. Moreover, physical interpretations can be attached to the associated CS. For instance, $SU(2)$ -CS [13] allow a semi-classical description of spin and the $SU(1, 1)$ -CS [12] are optimally localized states for the quantum mechanics of a free particle on the $(1 + 1)$ -dimensional anti-de Sitter spacetime. It is worth mentioning that the square integrability condition has recently been replaced by the less restrictive notion of square integrability modulo a subgroup, which allowed the construction of (quasi-)coherent states for the Poincaré group in $(1 + 1)$ -dimensions [14]. A more general framework is described in [15].

1.4. Let us now discuss the relevance of the CS to the quantization-versus-classical-limit procedures. A classical G -elementary system is generally described by a coadjoint orbit of G , which is a symplectic G -homogeneous space $(G/H, \omega)$, where H is a closed subgroup of G , and ω is a G -invariant, closed and non-degenerate 2-form on G/H . On the other hand, a quantum G -elementary system is described by a pair $(U, \mathcal{H})$, where U is a Unitary Irreducible Representation (UIR) of G in a Hilbert space $\mathcal{H}$. Classical and quantum G -elementary systems are related to each other by on the one hand, the quantization methods, such as the Kirillov-Kostant-Souriau geometric quantization (also known as the orbits method) [16, 17, 18], or Berezin's quantization [6], or the deformation quantization [19], and on the other hand the classical limit procedures, such as the CS-inspired one described by Onofri [7]. Since at different stages of this work we will be dealing with these two kinds of procedures, we now briefly hint at their intuitive content.

1.5. Whenever U is a square integrable UIR of G , one can construct a family of CS parametrized by G/H , where H is the closed subgroup of G which leaves invariant, up to a phase, the fiducial state. By construction, this family bears in its very structure enough information to allow one to equip G/H with a G -invariant symplectic form ω , which makes $(G/H, \omega)$ into the classical G -elementary system describing the classical limit of the quantum G -elementary system $(U, \mathcal{H})$. More precisely, this explicit construction leads to a Kähler G -homogeneous space [7]. Combining this derivation with the evaluation of Berezin's covariant symbols [6] provides one with a moment map that identifies $(G/H, \omega)$ with a coadjoint orbit of G . Berezin's symbols are the mean values of the quantum generators of $\mathfrak{g}$ in the CS.

1.6. Obviously, evaluating the classical limit is more natural than quantizing. However, because of their better understanding of classical theories, mathematical-physicists are very much interested in having a quantization procedure that would allow the translation of some of the well established classical physical understanding to a quantum counterpart. Several quantization procedures are now available. Three of them have already been mentioned. Geometric quantization is the method that we will be dealing with in this paper. In its simplest version, this technique associates a quantum G -elementary system to a classical one $(G/H, \omega)$, provided that $[\omega]$ is an integral cohomology class and that G/H admits an invariant polarization.

1.7. The very natural question we address in this work can be formulated as follows: *How do the quantization-versus-classical-limit procedures depicted above extend to the supersymmetric context?* We provide an answer to this, by studying very specific though non-trivial examples, namely those of the typical and the atypical $\mathrm{OSp}(2/2)$ -elementary systems. To shed more light on our motivations, we now situate our present contribution within the framework of the fast developing field of supermathematics.

1.8. Supermathematics is the collection of mathematical tools developed during the last thirty years in order to provide physicists with a rigorous framework for the study of the so-called supersymmetric theories, such as supergravity and superstrings. These are theories that possess symmetries which mix their bosonic and fermionic degrees of freedom. Some of the tools were already available before these theories really triggered the interest of a large number of researchers (see [20] pp. 26–28). A description of the super extensions of the usual analytic, algebraic and geometric concepts that have so far been obtained would unfortunately lead us too far from the subject of this paper; we refer the interested reader to the existing literature [21]–[31] and confine our description to those super ingredients that are crucial for answering the question raised in 1.7.

1.9. Contrary to Lie superalgebras, there exist several notions of Lie supergroups which correspond to the different definitions of a supermanifold (see [30] for more details). Concerning the representation theory, only abstract representations (Verma modules) of some simple Lie superalgebras are known [22, 24, 25, 32, 33]. Using these representations, supercoherent states have recently been explicitly constructed [34, 35, 36]. On the other hand, supergeometric concepts such as supersymplectic supermanifolds and super coadjoint orbits have been understood since the end of the 70s [22, 28] (see [37] for a recent application in physics). A very nice characterization of the latter in terms of usual geometric objects has been obtained by Rothstein [38] (see also [39]). Hence, all the ingredients needed for

answering the question formulated in 1.7 are available. It remains to super extend the methods described in 1.5 and 1.6. This is explicitly carried out here for the case of the typical and atypical $\mathrm{OSp}(2/2)$ -CS.

1.10. The paper is organized as follows: In Section 2 the so-called typical and atypical super unitary irreducible representations of the Lie superalgebra $\mathfrak{osp}(2/2)$, which super extend the discrete series representations of its subalgebra $\mathfrak{su}(1,1)$, are described. Then, the typical coherent states are constructed. By doing this we reproduce the construction of the $\mathrm{OSp}(2/2)$ -CS obtained in [35], and cure some of the discrepancies appearing there. In Section 3 a super extension of the methods depicted in 1.5 is then applied to these CS. Hence, the $\mathrm{OSp}(2/2)$ -homogeneous superspace parametrizing the latter, which we denote by $\mathcal{D}^{(1|2)}$, is explicitly equipped with an $\mathrm{OSp}(2/2)$ -invariant supersymplectic structure ω . Moreover, the evaluation of Berezin's covariant symbols exhibits a momentum map that identifies $(\mathcal{D}^{(1|2)}, \omega)$ with an $\mathrm{OSp}(2/2)$ -coadjoint orbit. Finally, the invariant super measure on $\mathcal{D}^{(1|2)}$ obtained in [35] is recovered through a simpler computation. In Section 4, after an introduction to the general theory, the supergeometry of $(\mathcal{D}^{(1|2)}, \omega)$ is studied further. As a result, $(\mathcal{D}^{(1|2)}, \omega)$ is shown to be not only a non-trivial example of Rothstein's general supersymplectic supermanifolds [38] but also a non-trivial example of the notion of a super Kähler supermanifold already discussed in [8, 9]. In this particular setting, we show how Rothstein's characterization can be improved. In Section 5, after a brief introduction to the usual geometric quantization procedure, a super extension of it is applied to $(\mathcal{D}^{(1|2)}, \omega)$. General superprequantization has been developed by Kostant [28], however the lack of a notion of polarization prevented him from completing the quantization. Here, the super Kähler character of $(\mathcal{D}^{(1|2)}, \omega)$ singles out a natural invariant super Kähler polarization which leads to the complete quantization. The coherent states associated to the atypical representations (atypical CS), their underlying supergeometry, and its quantization are described in Section 6. In particular the atypical coadjoint orbit is identified. Section 7 gathers additional results and discussions, while concluding remarks and possible extensions of the present work are displayed in Section 8. Our conventions and notations are presented in Appendix A, while useful constructions are relegated to Appendix B.

1.11. Before presenting the details, we would like to make clear some important points concerning our approach and strategy. Perelomov's construction of G -CS can be explicitly carried out starting simply from an infinitesimal version of a UIR of G [3]. This approach, used in [34, 35], is also adopted here. It might also seem strange that we start with a

representation of the Lie superalgebra, evaluate its classical limit through the associated CS and then quantize the obtained supersymplectic supermanifold in order to construct a representation of the same Lie superalgebra! In fact, the representation we start with is, as we will see, an abstract representation (Verma module), while the second is an explicit one. The latter, is realized in a super Hilbert space of superholomorphic sections of a line bundle sheaf over the considered coadjoint orbit. This representation is an important step towards constructing explicit super UIR of Lie supergroups.

2 $\mathfrak{osp}(2/2)$ representations and typical $\mathrm{OSp}(2/2)$ -CS

This section is devoted to two main purposes. We first describe the $\mathfrak{osp}(2/2)$ Lie superalgebra and its lowest weight typical and atypical representations, then we construct the associated coherent states.

2.1 $\mathfrak{osp}(2/2)$ representations

The superalgebra we consider here is the real orthosymplectic superalgebra $\mathfrak{osp}(2/2, \mathbb{R})$. Throughout, we will simply denote it $\mathfrak{osp}(2/2)$. It is a real non-compact form of the basic classical simple Lie superalgebra $\mathfrak{osp}(2/2, \mathbb{C})$, which corresponds to $C(2)$ in Kac's classification [25]. Its $\mathbb{Z}_2$ -grading, $\mathfrak{osp}(2/2) = \mathfrak{osp}(2/2)_{\bar{0}} \oplus \mathfrak{osp}(2/2)_{\bar{1}}$, is such that the even component $\mathfrak{osp}(2/2)_{\bar{0}} = \mathfrak{so}(2) \oplus \mathfrak{sp}(2, \mathbb{R})$, and the odd component $\mathfrak{osp}(2/2)_{\bar{1}}$ is an $\mathfrak{osp}(2/2)_{\bar{0}}$ -module. In what follows, using the isomorphism $\mathfrak{sp}(2, \mathbb{R}) \cong \mathfrak{su}(1, 1)$, we will consider $\mathfrak{su}(1, 1)$ instead of $\mathfrak{sp}(2, \mathbb{R})$.

As for Lie algebras, the construction of representations of $\mathfrak{osp}(2/2)$ relies on its complexification $\mathfrak{osp}(2/2, \mathbb{C})$. We now display a description of the latter superalgebra. Since $\mathfrak{osp}(2/2, \mathbb{C})$ is of type I [25, 23], its odd component $\mathfrak{osp}(2/2, \mathbb{C})_{\bar{1}}$ decomposes into two irreducible $\mathfrak{osp}(2/2, \mathbb{C})_{\bar{0}}$ -modules, namely, $\mathfrak{osp}(2/2, \mathbb{C})_{\bar{1}} = \mathfrak{osp}(2/2, \mathbb{C})_{-1} \oplus \mathfrak{osp}(2/2, \mathbb{C})_1$. Moreover $\mathfrak{osp}(2/2, \mathbb{C})$ admits the following $\mathbb{Z}_2$ -compatible $\mathbb{Z}$ -gradation:

$$\mathfrak{osp}(2/2, \mathbb{C}) = \mathfrak{osp}(2/2, \mathbb{C})_{-1} \oplus \mathfrak{osp}(2/2, \mathbb{C})_0 \oplus \mathfrak{osp}(2/2, \mathbb{C})_1, \quad (2.1)$$

where $\mathfrak{osp}(2/2, \mathbb{C})_0 = \mathfrak{osp}(2/2, \mathbb{C})_{\bar{0}}$. For $\{B, K_0, K_{\pm}; V_{\pm}, W_{\pm}\}$ a basis of $\mathfrak{osp}(2/2, \mathbb{C})$, the above structures are explicitly displayed in the following commutation ($[\cdot, \cdot]$) and anticom-

mutation ($[\cdot, \cdot]_+$) relations:

$$[K_0, K_\pm] = \pm K_\pm, \quad [K_+, K_-] = -2K_0, \quad (2.2a)$$

$$[B, K_\pm] = 0, \quad [B, K_0] = 0, \quad (2.2b)$$

$$[K_0, V_\pm] = \pm \frac{1}{2} V_\pm, \quad [K_0, W_\pm] = \pm \frac{1}{2} W_\pm, \quad (2.2c)$$

$$[K_\pm, V_\pm] = 0, \quad [K_\pm, W_\pm] = 0, \quad (2.2d)$$

$$[K_\pm, V_\mp] = \mp V_\pm, \quad [K_\pm, W_\mp] = \mp W_\pm, \quad (2.2e)$$

$$[B, V_\pm] = \frac{1}{2} V_\pm, \quad [B, W_\pm] = -\frac{1}{2} W_\pm, \quad (2.2f)$$

$$[V_\pm, V_\pm]_+ = 0, \quad [W_\pm, W_\pm]_+ = 0, \quad (2.2g)$$

$$[V_\pm, V_\mp]_+ = 0, \quad [W_\pm, W_\mp]_+ = 0, \quad (2.2h)$$

$$[V_\pm, W_\pm]_+ = K_\pm, \quad [V_\pm, W_\mp]_+ = K_0 \mp B. \quad (2.2i)$$

Clearly, the even component is spanned by $\{B, K_0, K_\pm\}$. The K -generators form an $\mathfrak{su}(1, 1)$ Lie subalgebra (see (2.2a)), and B spans a one dimensional center of $\mathfrak{osp}(2/2, \mathbb{C})_{\bar{0}}$ (see (2.2b)). On the other hand, the odd component is the span of $V_\pm$ and $W_\pm$. More precisely, $\{V_\pm\}$ (resp. $\{W_\pm\}$) span $\mathfrak{osp}(2/2, \mathbb{C})_1$ (resp. $\mathfrak{osp}(2/2, \mathbb{C})_{-1}$). The fact that each of these two-dimensional vector spaces carry an irreducible representation of $\mathfrak{osp}(2/2, \mathbb{C})_{\bar{0}}$ is transparent from equations (2.2c)–(2.2f); $\mathfrak{osp}(2/2, \mathbb{C})_1$ and $\mathfrak{osp}(2/2, \mathbb{C})_{-1}$ are distinguished by the distinct eigenvalues of B in (2.2f).

In order to construct irreducible highest (or lowest) weight representations of $\mathfrak{osp}(2/2)$ one needs to exhibit a Borel subsuperalgebra. In other words, we look for a decomposition of $\mathfrak{osp}(2/2, \mathbb{C})$ of the following form:

$$\mathfrak{osp}(2/2, \mathbb{C}) = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+, \quad (2.3)$$

where $\mathfrak{h}$ is a Cartan subalgebra of $\mathfrak{osp}(2/2, \mathbb{C})_{\bar{0}}$, and $\mathfrak{n}^-$ and $\mathfrak{n}^+$ are subsuperalgebras of $\mathfrak{osp}(2/2, \mathbb{C})$ such that $[\mathfrak{h}, \mathfrak{n}^\pm] \subseteq \mathfrak{n}^\pm$. The subsuperalgebras $\mathfrak{b}^+ = \mathfrak{h} \oplus \mathfrak{n}^+$ and $\mathfrak{b}^- = \mathfrak{h} \oplus \mathfrak{n}^-$ are called respectively the positive and the negative Borel subsuperalgebras. In terms of the basis of $\mathfrak{osp}(2/2, \mathbb{C})$ given above it clearly appears that $\mathfrak{h}$ is the complex span of $\{B, K_0\}$. Moreover, $\mathfrak{n}_0^+$ (resp. $\mathfrak{n}_0^-$) can be taken to be the span of $\{K_+\}$ (resp. $\{K_-\}$). Having fixed this, it can easily be shown that there exist three possible Borel subsuperalgebras of $\mathfrak{osp}(2/2, \mathbb{C})$. Since $\mathfrak{b}^\pm = \mathfrak{h} \oplus \mathfrak{n}_0^\pm \oplus \mathfrak{n}_1^\pm$, we need only to exhibit the three possible $\mathfrak{n}_1^\pm$. These are:

(i) $\mathfrak{n}_1^+ = \text{span}\{V_+, V_-\}$ and $\mathfrak{n}_1^+ = \text{span}\{W_+, W_-\}$,

(ii) $\mathfrak{n}_1^+ = \text{span}\{W_+, W_-\}$ and $\mathfrak{n}_1^- = \text{span}\{V_+, V_-\}$,

(iii) $\mathfrak{n}_1^+ = \text{span}\{V_+, W_+\}$ and $\mathfrak{n}_1^- = \text{span}\{V_-, W_-\}$.

Few remarks are now in order.

Remark 2.1: The situation in (i) is symmetric to that in (ii). Moreover, they both fit with the $\mathbb{Z}$ -grading given in (2.1). Indeed, we have that $\text{osp}(2/2, \mathbb{C})_0 = \mathfrak{n}_0^- \oplus \mathfrak{h} \oplus \mathfrak{n}_0^+$ and $\text{osp}(2/2, \mathbb{C})_{\pm 1} = \mathfrak{n}_1^{\pm}$.

Remark 2.2: The $\pm$ indices carried by the V s and the W s are misleading regarding the root space decomposition of $\text{osp}(2/2, \mathbb{C})$ relative to the $\mathfrak{h}$ given above. Indeed, one can see from (2.2i) that if V_+ (resp. W_+) is associated to some odd root α , then W_- (resp. V_-) is associated to $-\alpha$. This unconventional choice of notations is aimed at making the $\mathbb{Z}$ -gradation of (2.1) explicit in the defining relations of $\text{osp}(2/2, \mathbb{C})$ (see (2.2a)–(2.2i)).

Remark 2.3: Since the set of positive roots depends on the positive Borel subsuperalgebra considered, we have then here three possible sets of this type. While the set of positive roots arising from the case (iii) contains two odd simple roots, those arising from the cases (i) and (ii) contain one odd and one even simple roots (recall that $\text{rank}(\text{osp}(2/2, \mathbb{C})) = 2$). The systems of simple roots of both (i) and (ii) can be connected to each other by an element of the Weyl group of $\text{osp}(2/2, \mathbb{C})$ ($\mathcal{W}(\text{osp}(2/2, \mathbb{C})) := \mathcal{W}(\text{osp}(2/2, \mathbb{C})_{\bar{0}})$); they then give rise to the same Dynkin diagram. On the other hand, the system of simple roots of (iii) does not belong to the $\mathcal{W}(\text{osp}(2/2, \mathbb{C}))$ -orbit containing the two previous cases; a different Dynkin diagram arises then. This situation is a special feature of basic classical simple Lie superalgebras [40]. Indeed, usual complex simple Lie algebras admit a unique Dynkin diagram. The two Dynkin diagrams mentioned above exhaust all the possibilities for $\text{osp}(2/2, \mathbb{C})$ [40]. Notice finally that Kac considered in his original classification [25] only those Borel subsuperalgebras that lead to the minimum number of odd simple roots. They were called *distinguished* Borel subsuperalgebras.

Remark 2.4: Contrary to the two other cases, the choice in (iii) leads to a very interesting $\mathbb{Z}$ -grading of $\text{osp}(2/2, \mathbb{C})$ [33]. Both different and finer than that of equation (2.1), it is given by:

$$\text{osp}(2/2, \mathbb{C}) = \overset{(-2)}{\mathfrak{n}_0^-} \oplus \overset{(-1)}{\mathfrak{n}_1^-} \oplus \overset{(0)}{\mathfrak{h}} \oplus \overset{(1)}{\mathfrak{n}_1^+} \oplus \overset{(2)}{\mathfrak{n}_0^+}. \quad (2.4)$$

The abstract lowest weight representations that will be described below are those obtained using the Borel subsuperalgebras of case **(iii)**. This choice is justified simply by the fact that, up to some discrepancies that we correct here, the associated representations are those already used in [35]. A more mathematical description of these representations is given in [33].

An *abstract irreducible lowest-weight $osp(2/2)$ -module* is explicitly constructed starting from a lowest-weight state. According to our choice of Borel subsuperalgebras, namely the one made above in **(iii)**, the lowest-weight state is the state, temporarily denoted $|0\rangle$, which is simultaneously annihilated by K_- , V_- and W_- , and which moreover is a common eigenstate of both the Cartan subalgebra generators, B and K_0 . Hence, $|0\rangle$ is such that,

$$K_0|0\rangle = \tau|0\rangle, \quad B|0\rangle = b|0\rangle, \quad (2.5)$$

and

$$K_-|0\rangle = V_-|0\rangle = W_-|0\rangle = 0, \quad (2.6)$$

where $0 < \tau \in \mathbb{R}$ and $b \in \mathbb{R}$ completely specify a lowest-weight $osp(2/2)$ -module, which is denoted $V(\tau, b)$ throughout. A basis of $V(\tau, b)$ is explicitly obtained by applying to $|0\rangle$ basis elements of the enveloping superalgebra of the Lie subsuperalgebra $\mathfrak{n}^+$ (as given in (2.3) and **(iii)** above). The results of this construction are now displayed; more details are given in Appendix B.

The following observation simplifies the construction. By restricting (2.5) and (2.6) to the $\mathfrak{su}(1, 1)$ -generators, K_0 and K_- , it clearly appears that $|0\rangle$ is also the lowest-weight vector of an irreducible $\mathfrak{su}(1, 1)$ -module $D(\tau) \subset V(\tau, b)$, namely,

$$D(\tau) \equiv \text{span}\{|\tau, \tau + m\rangle, m \in \mathbb{N}\}. \quad (2.7)$$

This is the representation space of the well known *positive discrete series representations* of $\mathfrak{su}(1, 1)$. As a subspace of $V(\tau, b)$, $D(\tau)$ is an eigenspace of B with eigenvalue b . This is an immediate consequence of both (2.5) and the fact that B commutes with $\mathfrak{su}(1, 1)$ (see (2.2b)). Combined with (2.7), these facts suggest the following notation for the lowest-weight state of $V(\tau, b)$, namely $|0\rangle \equiv |b, \tau, \tau\rangle$.

As it is explicitly shown in Appendix B, $V(\tau, b)$ is built out of more than one irreducible $\mathfrak{su}(1, 1)$ -module. Two cases must however be distinguished, namely, either $|b| < \tau$ or $b = \pm\tau$.

Typical: When $|b| < \tau$, as a vector space, $V(\tau, b)$, turns out to be the direct sum of irreducible lowest-weight $\mathfrak{su}(1, 1)$ -modules (positive discrete series). More precisely,

$$V(\tau, b) \equiv D(\tau) \oplus 2 \cdot D(\tau + \tfrac{1}{2}) \oplus D(\tau + 1). \quad (2.8)$$

Here $D(\tau + \frac{1}{2})$ appears with multiplicity 2. These two copies of $D(\tau + \frac{1}{2})$ are distinguished eigenspaces of B , with eigenvalues $b + \frac{1}{2}$ and $b - \frac{1}{2}$. The degeneracy is then raised when one considers the $\mathfrak{su}(1, 1)$ -modules appearing in (2.8) as $(\mathfrak{su}(1, 1) \oplus \mathfrak{so}(2))$ -modules. An extra subscript has then to be added to our previous notation in order to take this fact into account. Observing that $D(\tau + 1)$, as $D(\tau)$, is a B -eigenspace with eigenvalue b , we can write the following finer decomposition of $V(\tau, b)$,

$$V(\tau, b) = D_b(\tau) \oplus D_{b+\frac{1}{2}}(\tau + \tfrac{1}{2}) \oplus D_{b-\frac{1}{2}}(\tau + \tfrac{1}{2}) \oplus D_b(\tau + 1). \quad (2.9)$$

More precisely,

$$V(\tau, b) \equiv \text{span}\{|b, \tau, \tau + m\rangle, |b + \tfrac{1}{2}, \tau + \tfrac{1}{2}, \tau + \tfrac{1}{2} + m\rangle, \\ |b - \tfrac{1}{2}, \tau + \tfrac{1}{2}, \tau + \tfrac{1}{2} + m\rangle, |b, \tau + 1, \tau + 1 + m\rangle; m \in \mathbb{N}\}. \quad (2.10)$$

Moreover, as a vector superspace, $V(\tau, b) = V_0(\tau, b) \oplus V_1(\tau, b)$, where $V_0(\tau, b) \equiv D_b(\tau) \oplus D_b(\tau + 1)$ and $V_1(\tau, b) \equiv D_{b+\frac{1}{2}}(\tau + \frac{1}{2}) \oplus D_{b-\frac{1}{2}}(\tau + \frac{1}{2})$. (The action of the generators of $\mathfrak{osp}(2/2, \mathbb{C})$ in $V(\tau, b)$ is explicitly given in Appendix B.) The irreducible $\mathfrak{osp}(2/2)$ -modules, $V(\tau, b)$, for $|b| < \tau$ are usually called *typical representations* [23, 24, 25, 32].

Atypical: When $b = \tau$ (resp. $b = -\tau$), $V(\tau, \tau)$ (resp. $V(\tau, -\tau)$) is no longer irreducible. It contains an $\mathfrak{osp}(2/2)$ -submodule, $V'(\tau, \tau) \equiv D_{\tau+\frac{1}{2}}(\tau + \frac{1}{2}) \oplus D_\tau(\tau + 1)$ (resp. $V'(\tau, -\tau) \equiv D_{-\tau-\frac{1}{2}}(\tau + \frac{1}{2}) \oplus D_{-\tau}(\tau + 1)$), generated by the *primitive vector* $V_+|0\rangle$ (resp. $W_+|0\rangle$). The quotient $V(\tau, \tau)/V'(\tau, \tau)$ (resp. $V(\tau, -\tau)/V'(\tau, -\tau)$) appears then as the appropriate irreducible $\mathfrak{osp}(2/2)$ -module. More precisely,

$$U(\pm\tau) \equiv V(\tau, \pm\tau)/V'(\tau, \pm\tau) \equiv D_{\pm\tau}(\tau) \oplus D_{\pm(\tau-\frac{1}{2})}(\tau + \tfrac{1}{2}). \quad (2.11)$$

These representations are known as the *atypical representations* [23, 24, 25, 32].

The above typical and atypical $\mathfrak{osp}(2/2)$ -modules can be turned into *super unitary* irreducible representations [33], by equipping $V(\tau, b)$ with a *super Hermitian* form $\langle \cdot | \cdot \rangle$.

The latter is a notion originally defined in [41], and used in the context of representation theory of Lie superalgebras in [24], and more recently in [33]. It is defined in the following way:

$$\langle \cdot | \cdot \rangle : V(\tau, b) \times V(\tau, b) \longrightarrow \mathbb{C} \quad (2.12)$$

such that $\forall u, v$ two homogeneous elements of $V(\tau, b)$

$$\overline{\langle u | v \rangle} = (-1)^{\epsilon(u)\epsilon(v)} \langle v | u \rangle, \quad (2.13)$$

where the parity of a homogeneous element $w \in V(\tau, b)$ is $\epsilon(w) \equiv 0(1)$ for $w \in V_{\bar{0}(\bar{1})}(\tau, b)$. The elements of $V_{\bar{0}(\bar{1})}$ are called even (odd) elements of $V(\tau, b)$. The super Hermitian form is taken linear in the second component. In what follows we will consider a *homogeneous* realization of (2.13), namely, $\forall u = u_0 + u_1, v = v_0 + v_1 \in V(\tau, b) = V_{\bar{0}}(\tau, b) \oplus V_{\bar{1}}(\tau, b)$,

$$\langle u | v \rangle \equiv \langle u_0 | v_0 \rangle_0 + i \langle u_1 | v_1 \rangle_1, \quad (2.14)$$

where, $\langle \cdot | \cdot \rangle_0$ (resp. $\langle \cdot | \cdot \rangle_1$) is an Hermitian form, in the usual sense, on $V_{\bar{0}}(\tau, b)$ (resp. $V_{\bar{1}}(\tau, b)$). Since $\overline{\langle u_k | v_k \rangle_k} = \langle v_k | u_k \rangle_k$ for $k = 0, 1$, one clearly sees that (2.14) satisfies (2.13). The super Hermitian form in (2.14) is even.

Besides being an $\mathfrak{osp}(2/2)$ -module, $V(\tau, b)$ will be also considered as a left $\mathcal{B}$ -module, where $\mathcal{B} = \mathcal{B}_0 \oplus \mathcal{B}_1$ is a *complex Grassmann algebra* [22, 23]. The super Hermitian form (2.12)–(2.14) will then be extended to the *Grassmann envelope of the second kind* [22] $\tilde{V}(\tau, b)$ of $V(\tau, b)$,

$$\tilde{V}(\tau, b) \equiv (\mathcal{B} \otimes V(\tau, b))_0, \quad (2.15)$$

such that

$$\langle \cdot | \cdot \rangle : \tilde{V}(\tau, b) \times \tilde{V}(\tau, b) \longrightarrow \mathcal{B}_0 \quad (2.16)$$

From the next section on, $\mathcal{B}$ will assume a very specific form, as the exterior algebra over $\mathbb{C}^4$, namely, $\mathcal{B} \equiv \bigwedge \mathbb{C}^4$ (see Appendix A for more details about (2.15) and (2.16)).

Remark 2.5: The super Hermitian form in (2.12) and (2.14) is *positive definite* on $V(\tau, b)$ for $|b| < \tau$, in the sense that the Hermitian forms on both $V_{\bar{0}}(\tau, b)$ and $V_{\bar{1}}(\tau, b)$ are positive definite. In Appendix B, we show how this structure turns the typical module $V(\tau, b)$ into a super unitary irreducible representation of $\mathfrak{osp}(2/2)$.

Remark 2.6: The construction of atypical representations (2.11) looks very much like the construction of the so-called indecomposable representations, which usually describe

massless relativistic quantum elementary systems. This analogy is confirmed by the fact that the super Hermitian form (2.14) is no longer positive definite on $V(\tau, \pm\tau)$. Indeed, as it is shown in Appendix B, both $V'(\tau, \tau)$ and $V'(\tau, -\tau)$ are $\mathfrak{osp}(2/2)$ -submodules of *zero-norm* states, with respect to the super Hermitian form. However, notice that the atypical representations are, as the typical ones, simply expressed in terms of discrete series representations of $\mathfrak{su}(1, 1)$ (see (2.11)).

Remark 2.7: It is important to note that the atypical modules $U(\pm\tau)$ in (2.11) are irreducible $\mathfrak{osp}(1/2)$ -modules, where $\mathfrak{osp}(1/2)$ stands here for the Lie subsuperalgebra of $\mathfrak{osp}(2/2)$ whose Cartan-Weyl basis is $\{K_0, K_{\pm}, \frac{1}{\sqrt{2}}(V_{\pm} + W_{\pm})\}$. This interesting observation will be discussed further on in Section 6.

We now briefly discuss Schur's lemma. Since $\mathfrak{osp}(2/2, \mathbb{C})$ is of rank 2, the center of its enveloping superalgebra is generated by two $\mathfrak{osp}(2/2, \mathbb{C})$ -invariants, which are, respectively, quadratic (the Casimir) and cubic in the generators of $\mathfrak{osp}(2/2, \mathbb{C})$. For simplicity, we exhibit below only the explicit expression of the former, which we denote Q_2 . Hence,

$$Q_2 = C_2 - B^2 + K_0 - W_+V_- - V_+W_-, \quad (2.17)$$

where C_2 is the Casimir invariant of the $\mathfrak{su}(1, 1)$ subalgebra, namely,

$$C_2 = K_0^2 - K_0 - K_+K_-. \quad (2.18)$$

The irreducibility of $V(\tau, b)$ implies that, on $V(\tau, b)$, both invariants are constant multiples of the identity. The exact value of the quadratic constant on $V(\tau, b)$ is simply obtained by evaluating Q_2 on the lowest-weight state $|b, \tau, \tau\rangle$. Hence,

$$Q_2 \equiv (\tau^2 - b^2)\mathbb{I} \quad \text{on} \quad V(\tau, b), \quad \text{since} \quad C_2 \equiv \tau(\tau - 1)\mathbb{I} \quad \text{on} \quad D(\tau) \subset V(\tau, b). \quad (2.19)$$

One clearly sees that Q_2 is identically zero on the atypical modules.

Finally, a lowest-weight irreducible $\mathfrak{osp}(2/2)$ -module is said to be an *integrable module*, whenever it is also a module for the *double covering* of the body $\mathrm{Sp}(2, \mathbb{R}) \times \mathrm{SO}(2)$ of $\mathrm{OSp}(2/2)$. Since the maximal torus in $\mathrm{OSp}(2/2)$ (and in $\mathrm{Sp}(2, \mathbb{R}) \times \mathrm{SO}(2)$ too) is the $U(1) \times U(1)$ subgroup generated by K_0 and B , one easily sees from (2.5) that $V(\tau, b)$ is an integrable $\mathfrak{osp}(2/2)$ -module if and only if both τ and b are half integers. Throughout we will only deal with integrable super unitary irreducible $\mathfrak{osp}(2/2)$ -modules, and we will no longer mention this fact. More precisely, we will restrict our attention to the typical

integrable representations, knowing that all the constructions of the forthcoming sections carry over to the atypical case. However, some very interesting points concerning the latter are worth to be mentioned. They are gathered in Section 6.

2.2 $\text{OSp}(2/2)$ coherent states

In this section we construct the $\text{OSp}(2/2)$ -coherent states associated to the typical representations described in the previous section. We should mention that, up to some discrepancies, that are cured here, and a different choice of conventions (see Appendix A), this construction was originally carried out in [35].

As stressed in the introduction, the coherent states for a Lie group G are the quantum states belonging to the orbit of G through a fiducial vector in an irreducible $\mathfrak{g}$ -module, carrying a unitary representation of G . Here, $\mathfrak{g}$ stands for the Lie algebra of G . Recall that we also mentioned in the introduction that the minimal requirement for the construction of G -CS consists in an irreducible unitary $\mathfrak{g}$ -module. For $\text{OSp}(2/2)$, irreducible super unitary $\text{osp}(2/2)$ -modules are at our disposal.

Considering the lowest-weight state $|b, \tau, \tau\rangle \in V(\tau, b)$ as the fiducial state, taking into account equations (2.5)–(2.6), and extending in a straightforward manner Perelomov's construction [3], the typical $\text{OSp}(2/2)$ -coherent states, are obtained as follows:

$$|a, \theta, \chi\rangle \equiv \mathcal{N} \exp(aK_+ + \theta V_+ + \chi W_+) |b, \tau, \tau\rangle. \quad (2.20)$$

They belong to $\tilde{V}(\tau, b)$ (see (2.10) and (2.15)), for $\mathcal{B}$ the Grassmann algebra generated by the complex anticommuting variables θ, χ , and their complex conjugates, $\bar{\theta}$ and $\bar{\chi}$. These are odd elements of $\mathcal{B}$, while a is an even element of $\mathcal{B}$. More details about $\mathcal{B}$ are given in Appendix A. In (2.20), $\mathcal{N}$ is a normalization factor which will be explicitly determined below.

Since, $[K_+, V_+] = 0$, $[K_+, W_+] = 0$ and $\theta^2 = \chi^2 = 0$, we can rewrite (2.20) as follows,

$$|a, \theta, \chi\rangle = \mathcal{N} [\exp(aK_+)] [1 + \theta V_+ + \chi W_+ + \frac{1}{2} \chi \theta (V_+ W_+ - W_+ V_+)] |b, \tau, \tau\rangle. \quad (2.21)$$

A simple computation, based on (B.15)–(B.19), leads to:

$$|a, \theta, \chi\rangle = \mathcal{N} \left[\exp \left(\left(a + \frac{b}{2\tau} \chi \theta \right) K_+ \right) \right] \left(|b, \tau, \tau\rangle + \theta \sqrt{\tau - b} |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2}\rangle \right. \\ \left. + \chi \sqrt{\tau + b} |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2}\rangle + \chi \theta \sqrt{\frac{(\tau^2 - b^2)(2\tau + 1)}{2\tau}} |b, \tau + 1, \tau + 1\rangle \right). \quad (2.22)$$

At this point, we introduce the new variable

$$z = a + \frac{b}{2\tau}\chi\theta \in \mathbb{C}. \quad (2.23)$$

Notice that for simplicity we are choosing here (and throughout) z to be a complex number. This choice is discussed in Section 7. Finally, using the known action of K_+ in $V(\tau, b)$ (B.18), we get the explicit form of the typical coherent states (2.22):

$$\begin{aligned} |z, \theta, \chi\rangle = \mathcal{N} & \left[\sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m)}{m! \Gamma(2\tau)}} z^m |b, \tau, \tau+m\rangle \right. \\ & + \theta \sqrt{\tau-b} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+1)}{m! \Gamma(2\tau+1)}} z^m |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle \\ & + \chi \sqrt{\tau+b} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+1)}{m! \Gamma(2\tau+1)}} z^m |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle \\ & \left. + \chi\theta \sqrt{\frac{(\tau^2-b^2)(2\tau+1)}{2\tau}} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+2)}{m! \Gamma(2\tau+2)}} z^m |b, \tau+1, \tau+1+m\rangle \right]. \end{aligned} \quad (2.24)$$

Now, using the super Hermitian form (2.16), the fact that the basis of $V(\tau, b)$ in (2.10) is super-orthonormal with respect to (2.14) (see Appendix B), and the conventions of Appendix A, one easily evaluates $\mathcal{N}$. Hence,

$$\mathcal{N} = (1 - |z|^2)^\tau \left[1 + i \frac{\tau-b}{2} \frac{\bar{\theta}\theta}{1-|z|^2} + i \frac{\tau+b}{2} \frac{\bar{\chi}\chi}{1-|z|^2} - \frac{(\tau^2-b^2)(\tau-1)}{4\tau} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1-|z|^2)^2} \right]. \quad (2.25)$$

Notice here that for obvious reasons we have considered $\mathcal{N}$ real. Moreover, the following identity,

$$\frac{1}{(1-x)^s} = \sum_{m=0}^{\infty} \frac{\Gamma(s+m)}{m! \Gamma(s)} x^m, \quad \text{for } |x| < 1, \quad (2.26)$$

has been used in order to write the result (2.25) in a compact form. It is worth mentioning at this point that when one sets to zero the odd variables θ and χ , all the previous formulae reduce exactly to those of the $SU(1, 1)$ -CS [3]. In that case, and clearly here too, z spans the unit disc $\mathcal{D}^{(1)} = \{z \in \mathbb{C} ; |z| < 1\}$; $\mathcal{D}^{(1)} \equiv SU(1, 1)/U(1)$.

For a later use, we evaluate now the $|\bar{z}, \bar{\theta}, \bar{\chi}\rangle$ CS, where, $\bar{z}$, $\bar{\theta}$ and $\bar{\chi}$ are the complex conjugates of z , θ and χ . Straightforwardly,

$$|\bar{z}, \bar{\theta}, \bar{\chi}\rangle = \mathcal{N}' \left[\sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m)}{m! \Gamma(2\tau)}} \bar{z}^m |b, \tau, \tau+m\rangle \right]$$

$$\begin{aligned}
& + \bar{\theta}\sqrt{\tau-b} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+1)}{m!\Gamma(2\tau+1)}} \bar{z}^m |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle \\
& + \bar{\chi}\sqrt{\tau+b} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+1)}{m!\Gamma(2\tau+1)}} \bar{z}^m |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle \\
& + \bar{\chi}\bar{\theta}\sqrt{\frac{(\tau^2-b^2)(2\tau+1)}{2\tau}} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+2)}{m!\Gamma(2\tau+2)}} \bar{z}^m |b, \tau + 1, \tau + 1 + m\rangle \Big],
\end{aligned} \tag{2.27}$$

the normalizing constant being now

$$\mathcal{N}' = (1 - |z|^2)^\tau \left[1 - i \frac{\tau-b}{2} \frac{\bar{\theta}\theta}{1 - |z|^2} - i \frac{\tau+b}{2} \frac{\bar{\chi}\chi}{1 - |z|^2} - \frac{(\tau^2-b^2)(\tau-1)}{4\tau} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1 - |z|^2)^2} \right]. \tag{2.28}$$

Having the explicit form of the typical $\text{OSp}(2/2)$ -CS, we can now apply to them a super extension of Onofri's analysis [7] in order to reveal the (super)geometry underlying them, or equivalently, in order to evaluate the classical limit of the quantum theory described by the typical representation $V(\tau, b)$ of the previous subsection. This analysis is the main concern of the next two sections.

3 Supersymplectic geometry and the $\text{OSp}(2/2)$ -CS

In analogy with the non-super case [3], the results of the previous section suggest that the space parametrizing the typical $\text{OSp}(2/2)$ -CS is the $\text{OSp}(2/2)$ -homogeneous superspace $\text{OSp}(2/2)/(U(1) \times U(1))$, realized in terms of the coordinates (z, θ, χ) , where z parametrizes the unit disc $\mathcal{D}^{(1)} := \{z \in \mathbb{C}; |z| < 1\}$. This realization will be subsequently called the $N = 2$ super unit disc, and it will be denoted $\mathcal{D}^{(1|2)}$. We recall that the $N = 1$ super unit disc $\mathcal{D}^{(1|1)}$ was fully considered in [8, 9]. We will consider here $\mathcal{D}^{(1|2)}$ as a supermanifold, although its complete and precise geometric characterization will be only given in Section 4.

In Section 3.1, we carry on super extending Onofri's analysis [7], which only makes use of the explicit form of the CS. This analysis will provide us with a partial description of the geometric structure underlying the typical $\text{OSp}(2/2)$ -CS. Some of the results obtained here will be revisited in Section 4, in the light of the general theory of supermanifolds. Let us recall that in the case of a semi-simple Lie group, starting from the associated CS, Onofri's analysis allows one to equip the homogeneous space parametrizing these CS with an invariant symplectic form (i.e. a closed and non-degenerate 2-form), which is moreover Kähler. Analogously, $\mathcal{D}^{(1|2)}$ is equipped here with an invariant *supersymplectic* form, in

the sense of [22, 28, 38]. Moreover, as it will be discussed in Section 4, this form is *super Kähler* in a sense to be defined (see also [8, 9]). In Section 3.2, extending Berezin's notion of covariant symbols, we evaluate the classical observables (superfunctions on $\mathcal{D}^{(1|2)}$) associated to the infinitesimal action of $\text{OSp}(2/2)$ on $\mathcal{D}^{(1|2)}$. By doing this we exhibit a moment map that identifies $\mathcal{D}^{(1|2)}$ with an $\text{OSp}(2/2)$ -coadjoint orbit. In the process, the Hamiltonian vector superfields associated to the classical observables are computed. Finally, in Section 3.3 we evaluate the *Liouville super measure* on $\mathcal{D}^{(1|2)}$. This will be needed in Section 5.

3.1 The supersymplectic form

Onofri's analysis starts by evaluating a real function from the G -CS; it plays the role of a Kähler potential for a G -invariant Kähler form on the space parametrizing the G -CS. Here, emphasizing the symplectic output of this procedure, we extend it to our typical $\text{OSp}(2/2)$ -CS. Hence, from equation (2.27), we evaluate the following superfunction on $\mathcal{D}^{(1|2)}$,

$$\begin{aligned} f(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi}) &:= \log |\langle 0 | \bar{z}, \bar{\theta}, \bar{\chi} \rangle|^{-2} \\ &= 2\tau \left[-\log(1 - |z|^2) + i \frac{\tau - b}{2\tau} \frac{\bar{\theta}\theta}{1 - |z|^2} \right. \\ &\quad \left. + i \frac{\tau + b}{2\tau} \frac{\bar{\chi}\chi}{1 - |z|^2} - \frac{\tau^2 - b^2}{4\tau^2} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1 - |z|^2)^2} \right]. \end{aligned} \quad (3.1)$$

Let now d be the exterior derivative on $\mathcal{D}^{(1|2)}$ given by $d = \delta + \bar{\delta}$, where

$$\delta = dz \frac{\partial}{\partial z} + d\theta \frac{\partial}{\partial \theta} + d\chi \frac{\partial}{\partial \chi}, \quad \bar{\delta} = d\bar{z} \frac{\partial}{\partial \bar{z}} + d\bar{\theta} \frac{\partial}{\partial \bar{\theta}} + d\bar{\chi} \frac{\partial}{\partial \bar{\chi}}. \quad (3.2)$$

Notice that here $\epsilon(d) = 0$, i.e. d is an even quantity.

An even two-superform on $\mathcal{D}^{(1|2)}$ can now be obtained from the superfunction in (3.1) in the following way:

$$\omega \equiv -i\delta\bar{\delta}f = \omega_0 + \omega_2 + \omega_4, \quad (3.3)$$

where,

$$\begin{aligned} \omega_0 &= \frac{-2i\tau}{(1 - |z|^2)^2} dz d\bar{z}; \\ \omega_2 &= -(\tau - b) \left[\frac{1}{(1 - |z|^2)} d\theta d\bar{\theta} + \frac{1}{(1 - |z|^2)^2} [\theta\bar{z} dz d\bar{\theta} - \bar{\theta}z d\theta d\bar{z}] - \bar{\theta}\theta \frac{(1 + |z|^2)}{(1 - |z|^2)^3} dz d\bar{z} \right] \end{aligned} \quad (3.4)$$

$$- (\tau + b) \left[\frac{1}{(1 - |z|^2)} d\chi d\bar{\chi} + \frac{1}{(1 - |z|^2)^2} [\chi \bar{z} dz d\bar{\chi} - \bar{\chi} z d\chi d\bar{z}] - \bar{\chi} \chi \frac{(1 + |z|^2)}{(1 - |z|^2)^3} dz d\bar{z} \right]; \quad (3.5)$$

$$\omega_4 = -i \frac{\tau^2 - b^2}{2\tau(1 - |z|^2)^2} \left[-2 \frac{(1 + 2|z|^2)}{(1 - |z|^2)^2} \bar{\theta} \bar{\chi} \chi \theta dz d\bar{z} + \bar{\theta} \theta d\chi d\bar{\chi} + \bar{\chi} \chi d\theta d\bar{\theta} - \bar{\theta} \chi d\theta d\bar{\chi} - \bar{\chi} \theta d\chi d\bar{\theta} + \frac{2\bar{\chi}\chi}{(1 - |z|^2)} [\theta \bar{z} dz d\bar{\theta} - \bar{\theta} z d\theta d\bar{z}] + \frac{2\bar{\theta}\theta}{(1 - |z|^2)} [\chi \bar{z} dz d\bar{\chi} - \bar{\chi} z d\chi d\bar{z}] \right]. \quad (3.6)$$

This two-superform belongs to the exterior algebra on $\mathcal{D}^{(1|2)}$. The latter is a bi-graded $\mathbb{Z} \times \mathbb{Z}_2$ algebra [28], where the $\mathbb{Z}$ -gradation is the usual gradation of de Rham complexes, while the $\mathbb{Z}_2$ -gradation is the natural gradation accompanying supersymmetry (i.e. the $\mathbb{Z}_2$ -gradation of the Grassmann algebra $\mathcal{B}$). More precisely, for any two superforms β_1 and β_2 on $\mathcal{D}^{(1|2)}$, one has:

$$\beta_1 \beta_2 = (-1)^{a_1 a_2 + b_1 b_2} \beta_2 \beta_1, \quad (3.7)$$

where a_i (resp. b_i) is the degree of the superform β_i with respect to the $\mathbb{Z}$ (resp. $\mathbb{Z}_2$) gradation. Hence, in (3.6), $dz d\bar{z} = -d\bar{z} dz$ (this is the usual wedge product), $dz d\bar{\theta} = -d\bar{\theta} dz$ and $d\theta d\bar{\theta} = d\bar{\theta} d\theta$.

Using these conventions one can check by explicit calculations that ω is closed, i.e. $d\omega = 0$. In fact, ω is closed by construction. This is a direct consequence of (3.3). Indeed, since $d = \delta + \bar{\delta}$, $d^2 = 0$ implies that $\delta^2 = \bar{\delta}^2 = \delta\bar{\delta} + \bar{\delta}\delta = 0$. Hence, $(\mathcal{D}^{(1|2)}, \omega)$ is a supersymplectic supermanifold. This particular point will be analyzed further on in Section 4.

3.2 The classical observables

By analogy with the non-super case, one can determine the classical observables associated to the generators of the supersymplectic action of $\text{OSp}(2/2)$ on $\mathcal{D}^{(1|2)}$. This is achieved through the evaluation of the so-called Berezin covariant symbols [6]. As for the superfunction (3.1), these are obtained simply from the knowledge of the explicit form of the $\text{OSp}(2/2)$ -CS and the representation $V(\tau, b)$ they belong to. Hence, the classical observable H^{cl} associated to an $\text{OSp}(2/2)$ -generator $H \in \{B, K_0, K_{\pm}, V_{\pm}, W_{\pm}\}$ is given by the Berezin symbol:

$$H^{cl} \equiv H^{cl}(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi}) := \langle \bar{z}, \bar{\theta}, \bar{\chi} | H | \bar{z}, \bar{\theta}, \bar{\chi} \rangle. \quad (3.8)$$

After lengthy but straightforward computations, based on (2.27) and results from Appendix B, one obtains:

$$\begin{aligned}
B^{cl} &= b + i \frac{\tau - b}{2} \frac{\bar{\theta}\theta}{1 - |z|^2} - i \frac{\tau + b}{2} \frac{\bar{\chi}\chi}{1 - |z|^2}, \\
K_0^{cl} &= \tau \frac{1 + |z|^2}{1 - |z|^2} \left[1 + i \frac{\tau - b}{2\tau} \frac{\bar{\theta}\theta}{(1 - |z|^2)} + i \frac{\tau + b}{2\tau} \frac{\bar{\chi}\chi}{(1 - |z|^2)} - \frac{\tau^2 - b^2}{2\tau^2} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1 - |z|^2)^2} \right], \\
K_+^{cl} &= \frac{2\tau z}{1 - |z|^2} \left[1 + i \frac{\tau - b}{2\tau} \frac{\bar{\theta}\theta}{(1 - |z|^2)} + i \frac{\tau + b}{2\tau} \frac{\bar{\chi}\chi}{(1 - |z|^2)} - \frac{\tau^2 - b^2}{2\tau^2} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1 - |z|^2)^2} \right], \\
K_-^{cl} &= \frac{2\tau \bar{z}}{1 - |z|^2} \left[1 + i \frac{\tau - b}{2\tau} \frac{\bar{\theta}\theta}{(1 - |z|^2)} + i \frac{\tau + b}{2\tau} \frac{\bar{\chi}\chi}{(1 - |z|^2)} - \frac{\tau^2 - b^2}{2\tau^2} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1 - |z|^2)^2} \right], \\
V_+^{cl} &= \frac{1}{1 - |z|^2} \left[-i(\tau - b)\theta + (\tau + b)z\bar{\chi} - \frac{\tau^2 - b^2}{2\tau} \frac{(iz\bar{\theta} + \chi)\bar{\chi}\theta}{(1 - |z|^2)} \right], \\
W_-^{cl} &= \frac{1}{1 - |z|^2} \left[(\tau - b)\bar{\theta} - i(\tau + b)\bar{z}\chi - \frac{\tau^2 - b^2}{2\tau} \frac{(\bar{z}\theta + i\bar{\chi})\bar{\theta}\chi}{(1 - |z|^2)} \right], \\
V_-^{cl} &= \frac{1}{1 - |z|^2} \left[-i(\tau - b)\bar{z}\theta + (\tau + b)\bar{\chi} - \frac{\tau^2 - b^2}{2\tau} \frac{(\bar{z}\chi + i\bar{\theta})\bar{\chi}\theta}{(1 - |z|^2)} \right], \\
W_+^{cl} &= \frac{1}{1 - |z|^2} \left[(\tau - b)z\bar{\theta} - i(\tau + b)\chi - \frac{\tau^2 - b^2}{2\tau} \frac{(iz\bar{\chi} + \theta)\bar{\theta}\chi}{(1 - |z|^2)} \right].
\end{aligned} \tag{3.9}$$

The obtained classical observables satisfy the following relations:

$$\overline{B^{cl}} = B^{cl}, \quad \overline{K_0^{cl}} = K_0^{cl}, \quad \overline{K_+^{cl}} = K_-^{cl}, \quad \overline{V_+^{cl}} = iW_-^{cl}, \quad \overline{V_-^{cl}} = iW_+^{cl}. \tag{3.10}$$

Remark 3.1: Notice that the Berezin symbols (3.8) are defined in terms of $|\bar{z}, \bar{\theta}, \bar{\chi}\rangle$ instead of $|z, \theta, \chi\rangle$. In order to justify this choice we need to anticipate on future results. In fact, as it will be shown in Section 5, this choice leads to the classical theory, the quantization of which gives rise to a superholomorphic representation. The anti-superholomorphic one arises as the quantization of the classical theory obtained from the CS $|z, \theta, \chi\rangle$.

The Hamiltonian vector superfield X_H associated to a classical observable H^{cl} , is the solution of the following defining equation [28],

$$X_H \lrcorner \omega = dH^{cl}, \tag{3.11}$$

where ‘ $\lrcorner$ ’ stands for the inner product and ω is the supersymplectic form (3.3). Here we display the Hamiltonian vector fields associated to the above observables. A long computation leads to the following:

$$\begin{aligned}
X_B &= i\frac{\theta}{2}\frac{\partial}{\partial\theta} - i\frac{\bar{\theta}}{2}\frac{\partial}{\partial\bar{\theta}} - i\frac{\chi}{2}\frac{\partial}{\partial\chi} + i\frac{\bar{\chi}}{2}\frac{\partial}{\partial\bar{\chi}}, \\
X_{K_0} &= iz\frac{\partial}{\partial z} - i\bar{z}\frac{\partial}{\partial\bar{z}} + i\frac{\theta}{2}\frac{\partial}{\partial\theta} - i\frac{\bar{\theta}}{2}\frac{\partial}{\partial\bar{\theta}} + i\frac{\chi}{2}\frac{\partial}{\partial\chi} - i\frac{\bar{\chi}}{2}\frac{\partial}{\partial\bar{\chi}}, \\
X_{K_+} &= iz^2\frac{\partial}{\partial z} - i\frac{\partial}{\partial\bar{z}} + iz\theta\frac{\partial}{\partial\theta} + iz\chi\frac{\partial}{\partial\chi}, \\
X_{K_-} &= i\frac{\partial}{\partial z} - i\bar{z}^2\frac{\partial}{\partial\bar{z}} - i\bar{z}\bar{\theta}\frac{\partial}{\partial\bar{\theta}} - i\bar{z}\bar{\chi}\frac{\partial}{\partial\bar{\chi}}, \\
X_{V_+} &= \left(\frac{\tau-b}{2\tau}\right)z\theta\frac{\partial}{\partial z} - i\left(\frac{\tau+b}{2\tau}\right)\bar{\chi}\frac{\partial}{\partial\bar{z}} + i\frac{\partial}{\partial\bar{\theta}} - \left(z + \left(\frac{\tau-b}{2\tau}\right)\chi\theta\right)\frac{\partial}{\partial\chi}, \\
X_{W_-} &= \left(\frac{\tau+b}{2\tau}\right)\chi\frac{\partial}{\partial z} - i\left(\frac{\tau-b}{2\tau}\right)\bar{z}\bar{\theta}\frac{\partial}{\partial\bar{z}} - \frac{\partial}{\partial\theta} + i\left(\bar{z} - \left(\frac{\tau-b}{2\tau}\right)\bar{\theta}\bar{\chi}\right)\frac{\partial}{\partial\bar{\chi}}, \\
X_{V_-} &= \left(\frac{\tau-b}{2\tau}\right)\theta\frac{\partial}{\partial z} - i\left(\frac{\tau+b}{2\tau}\right)\bar{z}\bar{\chi}\frac{\partial}{\partial\bar{z}} + i\left(\bar{z} + \left(\frac{\tau+b}{2\tau}\right)\bar{\theta}\bar{\chi}\right)\frac{\partial}{\partial\bar{\theta}} - \frac{\partial}{\partial\chi}, \\
X_{W_+} &= \left(\frac{\tau+b}{2\tau}\right)z\chi\frac{\partial}{\partial z} - i\left(\frac{\tau-b}{2\tau}\right)\bar{\theta}\frac{\partial}{\partial\bar{z}} - \left(z - \left(\frac{\tau+b}{2\tau}\right)\chi\theta\right)\frac{\partial}{\partial\theta} + i\frac{\partial}{\partial\bar{\chi}}.
\end{aligned} \tag{3.12}$$

The above vector fields verify:

$$\bar{X}_B = X_B, \quad \bar{X}_{K_0} = X_{K_0}, \quad \bar{X}_{K_+} = X_{K_-}, \quad \bar{X}_{V_+} = iX_{W_-}, \quad \bar{X}_{V_-} = iX_{W_+}. \tag{3.13}$$

The supersymplectic structure ω (3.3) on $\mathcal{D}^{(1|2)}$ defines a Poisson super bracket structure, $\{, \}$, in the space of smooth superfunctions on $\mathcal{D}^{(1|2)}$, turning it into a Poisson superalgebra. Indeed, given any two smooth superfunctions g and h on $\mathcal{D}^{(1|2)}$,

$$\{g, h\} \equiv -iX_g \lrcorner dh. \tag{3.14}$$

A simple computation shows that the classical observables in (3.9) form a Poisson sub-superalgebra isomorphic to $\mathfrak{osp}(2/2)$. Hence, the classical observables provide one with a supersymplectic realization of $\mathfrak{osp}(2/2)$. What precedes is equivalent to say that by evaluating the classical observables in (3.9) we have in fact exhibited an infinitesimally equivariant momentum map [18], $J : \mathcal{D}^{(1|2)} \rightarrow \mathfrak{osp}(2/2)^* \equiv \mathfrak{osp}(2/2)$, that identifies $\mathcal{D}^{(1|2)}$ with an $\mathrm{OSp}(2/2)$ -coadjoint orbit [28]. The latter is realized as a $(2|4)$ -dimensional subsupermanifold of $\mathbb{R}^{(4|4)}$, defined by two constraints, which correspond to the quadratic and the cubic $\mathfrak{osp}(2/2)$ -invariants. Indeed, when the latter are evaluated in the supersymplectic realization of $\mathfrak{osp}(2/2)$, they are both identically constant. For instance, $Q_2^{cl} \equiv \tau^2 - b^2$, (recall here that $|b| < \tau$).

This completes the first stage of our description of the geometry underlying the typical $\text{OSp}(2/2)$ -CS. This turns out to be a supersymplectic geometry. The main purpose of the next and final stage (Section 4) is to situate the results of the present section within the already existing theory of supersymplectic supermanifolds. Moreover, by analogy with the non super case [7], one is tempted to go one step further and consider $(\mathcal{D}^{(1|2)}, \omega)$ as an example of a general notion of *super Kähler supermanifolds*, which has not so far been seriously studied (see however [8, 9]). In Section 4, this notion will be given a legitimacy, which will open the door to super extending the full geometric quantization to the super Kähler context (see Section 5).

Before carrying out this program, we display now a computation, the result of which will be needed later on in Section 5.

3.3 The Liouville supermeasure

An super measure on the super unit disc $\mathcal{D}^{(1|2)}$, which super extends the notion of a Liouville measure on a symplectic manifold, can now be evaluated starting from ω (3.3) and using Berezin's notion of a density [29]. Up to a multiplicative constant, an $\text{OSp}(2/2)$ -invariant measure on $\mathcal{D}^{(1|2)}$ is given by:

$$d\mu(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi}) = \frac{i}{\pi} \text{sdet} ||\omega_{A\bar{B}}|| \, dz \, d\bar{z} \, d\theta \, d\bar{\theta} \, d\chi \, d\bar{\chi} \quad (3.15)$$

where 'sdet' stands for the superdeterminant (or Berezinian) [22], while $||\omega_{A\bar{B}}||$ stands for the supermatrix form of ω , namely,

$$\begin{aligned} \omega &\equiv dx^A \omega_{A\bar{B}} dx^{\bar{B}} \\ &= dz(a)d\bar{z} + dz(\alpha)d\bar{\theta} + dz(\beta)d\bar{\chi} + d\theta(\gamma)d\bar{z} + d\chi(\delta)d\bar{z} \\ &\quad + d\theta(y)d\bar{\theta} + d\theta(r)d\bar{\chi} + d\chi(s)d\bar{\theta} + d\chi(t)d\bar{\chi}. \end{aligned} \quad (3.16)$$

More precisely,

$$||\omega_{A\bar{B}}|| = \begin{pmatrix} a & \alpha & \beta \\ \gamma & y & r \\ \delta & s & t \end{pmatrix}; \quad (3.17)$$

the entries of $||\omega_{A\bar{B}}||$ are as follows:

$$\begin{aligned} a &= -\frac{i}{x^2} \left[2\tau + i(\tau - b) \frac{1 + |z|^2}{x} \bar{\theta}\theta + i(\tau + b) \frac{1 + |z|^2}{x} \bar{\chi}\chi - \frac{\tau^2 - b^2}{\tau} \frac{1 + 2|z|^2}{x^2} \bar{\theta}\bar{\chi}\chi\theta \right]; \\ \alpha &= -\frac{1}{x^2} (\tau - b) \bar{z}\theta \left[1 + i \frac{\tau + b}{\tau} \frac{\bar{\chi}\chi}{x} \right]; \quad \gamma = -\frac{i}{x^2} (\tau - b) z\bar{\theta} \left[1 + i \frac{\tau + b}{\tau} \frac{\bar{\chi}\chi}{x} \right]; \end{aligned}$$

$$\begin{aligned}
\beta &= -\frac{1}{x^2}(\tau + b)\bar{z}\chi \left[1 + i\frac{\tau - b}{\tau}\frac{\bar{\theta}\theta}{x} \right]; & \delta &= -\frac{i}{x^2}(\tau + b)z\bar{\chi} \left[1 + i\frac{\tau - b}{\tau}\frac{\bar{\theta}\theta}{x} \right]; \\
y &= -\frac{i}{x}(\tau - b) \left[1 + i\frac{\tau + b}{2\tau}\frac{\bar{\chi}\chi}{x} \right]; & t &= -\frac{i}{x}(\tau + b) \left[1 + i\frac{\tau - b}{2\tau}\frac{\bar{\theta}\theta}{x} \right]; \\
r &= -\frac{\tau^2 - b^2}{2\tau x^2}\bar{\theta}\chi; & s &= -\frac{\tau^2 - b^2}{2\tau x^2}\bar{\chi}\theta; & x &= 1 - |z|^2.
\end{aligned} \tag{3.18}$$

A simple computation leads then to:

$$d\mu(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi}) = \frac{-2\tau}{\pi(\tau^2 - b^2)} dz d\bar{z} d\theta d\bar{\theta} d\chi d\bar{\chi}. \tag{3.19}$$

Clearly, this measure is only valid in the typical case, $|b| < \tau$. (For the atypical case see Section 6.) Moreover, the choice of normalization in (3.19) is not innocent. Its usefulness will appear in Section 5 (see Remark 5.4). The $\text{OSp}(2/2)$ -invariance of $d\mu$ is claimed without proof. In fact, we can show that this is true using the action of $\text{OSp}(2/2)$ on $\mathcal{D}^{(1|2)}$. The latter can be derived by integrating the flows of the Hamiltonian vector fields (3.12) (see Section 7, 7.5).

It is worth mentioning at this point that up to a slight adaptation of conventions (see Appendix A), the supermeasure (3.19) is exactly the one used in [35] in order to prove the resolution of the identity for the typical $\text{OSp}(2/2)$ -CS. The same result holds here. In our notation, this means that the typical $\text{OSp}(2/2)$ -CS (2.27) form an overcomplete basis of $V(\tau, b)$ (see Section 7, 7.1). From a computational point of view the way we evaluate here (3.19) is by far simpler than the one used in [35].

4 More about supergeometry

The theory of supermanifolds was originally introduced in order to provide physicists with a rigorous framework for studying supersymmetric field theories. Here we are interested in later developments of the theory that were oriented towards super extending such techniques as symplectic geometry and geometric quantization. We start then this section by presenting a brief account of the key contributions in that direction (Section 4.1). The results of Section 3 will then be rediscussed in the light of the general theory (Section 4.2 and 4.3).

It is worth stressing that the most important point of this section is the extension of Rothstein's characterization of supersymplectic supermanifolds to the Kähler context (Section 4.3). This extension was already discussed in [8, 9] where a definition and a

non-trivial example of the notion of a super Kähler supermanifold were exhibited. Here, that definition is made more precise, explicit general formulae are given, and another non-trivial example is discussed, namely $(\mathcal{D}^{(1|2)}, \omega)$.

4.1 Supersymplectic supermanifolds

It is known that the geometry of a manifold can be recovered from its structure sheaf, i.e. the algebra of functions on that manifold. A supermanifold is defined by extending such an algebra to a supercommutative superalgebra. The supergeometry of this supermanifold is then extracted from the superstructure sheaf thus obtained through known techniques [28]. As for usual manifolds, three types of supermanifolds emerge, namely, the C^∞ , the real-analytic and the complex-analytic (or holomorphic) supermanifolds. From the results of the previous section it clearly appears that we are dealing here with the last type of supermanifolds.

Definition 4.1: [42, 43] *A $(p|q)$ -dimensional holomorphic supermanifold is a pair $(M, \mathcal{A}_M)$, where M is a p -dimensional complex manifold with holomorphic structure sheaf $\mathcal{O}_M$ and $\mathcal{A}_M$ is a sheaf of supercommutative superalgebras on M , such that:*

- (a) $\mathcal{A}_M/\mathcal{N}$ is isomorphic to $\mathcal{O}_M$, $\mathcal{N}$ being the subsheaf of nilpotent elements of $\mathcal{A}_M$, and
- (b) $\mathcal{A}_M$ is locally isomorphic to the exterior sheaf $\bigwedge \mathcal{E}$, where $\mathcal{E} \equiv \mathcal{N}/\mathcal{N}^2$ is a locally free sheaf over $\mathcal{O}_M$; equivalently, for $\{U_\alpha\}$ an open cover of M , $\mathcal{A}_M(U_\alpha)$ is locally isomorphic to $\mathcal{O}_M(U_\alpha) \otimes \bigwedge \mathbb{C}^q \equiv \bigwedge \mathcal{E}(U_\alpha)$.

Here $\bigwedge \mathbb{C}^q$ stands for the exterior algebra on $\mathbb{C}^q$, and p (resp. q) is the even (resp. odd) complex dimension of $(M, \mathcal{A}_M)$. Moreover, local supercoordinates on $(M, \mathcal{A}_M)$ are given by a set $(z^1, \dots, z^p; \theta^1, \dots, \theta^q)$, where $(z^1, \dots, z^p)$ are local coordinates on M and $(\theta^1, \dots, \theta^q)$ form a basis of $\mathcal{E}$ over $\mathcal{O}_M$. Notice that up to obvious modifications, Definition 4.1. applies equally well to C^∞ and real-analytic supermanifolds.

It is well known that a holomorphic vector bundle over M is completely specified given its sheaf of holomorphic sections, which is a locally free sheaf over $\mathcal{O}_M$. Hence, $\mathcal{E}$ in Definition 4.1 represents the sheaf of sections of a rank- q holomorphic vector bundle $\mathbb{F}$ over M . To any holomorphic supermanifold $(M, \mathcal{A}_M)$ one can then canonically associate the holomorphic supermanifold $(M, \mathcal{O}_M(\bigwedge \mathbb{F}))$, where $\mathcal{O}_M(\bigwedge \mathbb{F})$ is the sheaf of sections of the exterior bundle $\bigwedge \mathbb{F} \rightarrow M$. Condition (b) above implies that $(M, \mathcal{A}_M)$ and $(M, \mathcal{O}_M(\bigwedge \mathbb{F}))$ are locally isomorphic. The same holds true for C^∞ and real-analytic supermanifolds.

In these two last instances the local isomorphism always extends to a global but non-canonical one [44]. However, this is not always true in the holomorphic case [42, 43].

As for usual manifolds, the tangent sheaf is defined as the sheaf of superderivations of $\mathcal{A}_M$, and the cotangent sheaf $\Omega^1(\mathcal{A}_M)$ as its dual. Then a super de Rham complex can be constructed by introducing a coboundary operator d . From these ingredients one defines the notion of a supersymplectic supermanifold.

Definition 4.2: [28] *A supersymplectic supermanifold is a triple $(M, \mathcal{A}_M, \omega)$, where ω is a closed and non-degenerate even 2-superform on $(M, \mathcal{A}_M)$.*

Rothstein's characterization of C^∞ supersymplectic supermanifolds in terms of the geometry of vector bundles over usual symplectic manifolds constitutes a major contribution to this topic [38]. Indeed, for ω *at most quadratic in the odd coordinates*, Rothstein's theorem [38] allows one to completely identify ω in terms of a symplectic structure on M and extra structures in the vector bundle sector. More precisely, using the global isomorphism mentioned above, it states that to any C^∞ supersymplectic supermanifold $(M, \mathcal{A}_M, \omega)$ there corresponds a set $(M, \omega_0, \mathbb{E}, g, \nabla^g)$, where (M, ω_0) is a symplectic manifold, $\mathbb{E}$ is a vector bundle over M with metric g and g -compatible connection ∇^g , such that $\mathcal{E}$ is the sheaf of linear functionals on $\mathbb{E}$ and ω is completely determined in terms of (ω_0, g, ∇^g) as follows:

$$\omega = \omega_0 - d\alpha_2, \quad \text{where} \quad \alpha_2 = -g_{ab}\theta^a D\theta^b. \quad (4.1)$$

Here D is an operator defined on $\bigwedge \mathcal{E}$, with values in $\Omega^1(\mathcal{A}_M)$, such that:

$$D\theta^a \equiv d\theta^a - A_{ib}^a \theta^b dx^i, \quad a, b \in \{1, \dots, q\}, i \in \{1, \dots, p = 2n\}, \quad (4.2)$$

where $(x^i; \theta^a)$ are now real supercoordinates on $(M, \mathcal{A}_M, \omega)$, and A_{ib}^a are the components of ∇^g in the basis of the generators θ^a of $\mathcal{E}$. The explicit form of (4.1) is given by:

$$\omega = \omega_0 + \frac{1}{2}g_{ab}R_{ijc}^b \theta^c \theta^a dx^i dx^j + g_{ab}D\theta^a D\theta^b, \quad (4.3)$$

where R_{ijc}^b are the components of $\text{curv} \nabla^g$.

The correspondence mentioned above is one to one only in the C^∞ case. In the complex-analytic case either one considers supersymplectic holomorphic supermanifolds in the form $(M, \mathcal{O}_M(\bigwedge \mathbb{F}), \omega)$, for $\mathbb{F}$ a holomorphic vector bundle over M , or uses only the one way correspondence of Rothstein's theorem [38]. In both situations, equations similar to (4.1)–(4.3) hold; they are explicitly derived in Section 4.3. In what follows, *Rothstein's*

data will refer to the set $(M, \omega_0, \mathbb{E}, g, \nabla^g)$ associated to a supersymplectic supermanifold $(M, \mathcal{A}_M, \omega)$.

Few precisions are now in order. To be able to decompose without any ambiguity an even two-superform as a sum of homogeneous components in the anticommuting variables, one needs to restrict bundle automorphisms of $\bigwedge^2 \mathcal{E}$ to those automorphisms induced from bundle automorphisms of $\mathcal{E}$ [38]. Moreover, when ω contains terms of higher order in the odd coordinates (more than quadratic), the second part of Rothstein's theorem [38] states that there exists a superdiffeomorphism ρ of $(M, \mathcal{A}_M)$ such that ρ is the identity modulo $\bigwedge^2 \mathcal{E}$, and $\rho^*(\omega)$ is at most quadratic in the odd coordinates. Hence, the first part of Rothstein's theorem can be applied to $\rho^*(\omega)$. In other words, for $\omega = \omega_0 + \omega_2 + \omega_4 + \dots$, where the subscripts refer to the degree of homogeneity in the odd coordinates, one needs first to find the superdiffeomorphism ρ (which depends only on $\omega_4 + \dots$), then one uses it in order to transform ω into a 2-superform at most quadratic in the odd coordinates. Then one can identify Rothstein's data for ω . On the other hand, given those data one cannot reconstruct the original 2-superform. Indeed, only the transformed one is at reach, since ρ cannot be deduced from the above data. The supersymplectic form considered here (3.3)–(3.6), is obviously of the preceding form, nevertheless we will show in Section 4.3 that ω_4 can be obtained explicitly from Rothstein's data (as $\omega_0 + \omega_2$), without any mention to ρ (see (4.18)). In other words, the whole supersymplectic form can be obtained from the simple knowledge of $(M, \omega_0, \mathbb{E}, g, \nabla^g)$. This seems to be a common feature of super coadjoint orbits of the type considered here.

Our task is now twofold. First, identify $\mathcal{D}^{(1|2)}$ as a holomorphic supermanifold, then identify Rothstein's data for the supersymplectic $(\mathcal{D}^{(1|2)}, \omega)$. The first part is straightforward. For the second one, we will make use of a very useful Lemma proved in [8] in the case of $\mathcal{D}^{(1|1)} \equiv \text{OSp}(1/2)/\text{U}(1)$, and which applies to more general situations, in particular to the one in hand.

4.2 $\mathcal{D}^{(1|2)}$ as a holomorphic supermanifold

As a holomorphic supermanifold, the $\text{OSp}(2/2)$ -homogeneous superspace $\mathcal{D}^{(1|2)}$ obtained in Section 3 corresponds to the pair $(\mathcal{D}^{(1)}, \mathcal{A}_{\mathcal{D}^{(1)}})$, where $\mathcal{D}^{(1)}$ is the unit disc, and $\mathcal{A}_{\mathcal{D}^{(1)}} \equiv \mathcal{A}^{(1|2)} = \mathcal{O}_{\mathcal{D}^{(1)}} \otimes \bigwedge^2 \mathbb{C}^2$ is the defining superstructure sheaf. A general section h of $\mathcal{A}^{(1|2)}$ is a superholomorphic function, i.e.

$$h(z, \theta, \chi) \equiv h_1(z) + \theta h_2(z) + \chi h_3(z) + \chi \theta h_4(z), \quad (4.4)$$

where the h_i 's are holomorphic functions on $\mathcal{D}^{(1)}$. Moreover, the vector bundle $\mathbb{F}$ is nothing but a trivial rank 2 holomorphic vector bundle over $\mathcal{D}^{(1)}$.

Finally, notice here that the super observables in (3.9) are sections of the complexified superstructure sheaf $\mathcal{A}_{\mathbb{C}}^{(1|2)} \equiv C_{\mathbb{C}}^{\infty} \otimes \bigwedge \mathbb{C}^4$; and the Hamiltonian vector superfields in (3.12) are derivations of $\mathcal{A}_{\mathbb{C}}^{(1|2)}$. The canonically associated vector bundle is $\mathbb{F} \oplus \overline{\mathbb{F}}$, where $\overline{\mathbb{F}}$ is the complex conjugate bundle of $\mathbb{F}$.

4.3 $\mathcal{D}^{(1|2)}$ as a super Kähler supermanifold

Let us now identify the data $(M, \omega_0, \mathbb{E}, g, \nabla^g)$ for $(\mathcal{D}^{(1|2)}, \omega)$. First of all, it is straightforward from (3.6), that the symplectic manifold (M, ω_0) is here the symplectic unit disc $(\mathcal{D}^{(1)}, \omega_0)$, where $\omega_0 \equiv -2i\tau(1 - |z|^2)^{-2}dzd\bar{z}$. Indeed, observe that $\mathrm{SU}(1, 1) \times \mathrm{U}(1)$ is the body of $\mathrm{OSp}(2/2)$, and hence the body of $\mathcal{D}^{(1|2)} \equiv \mathrm{OSp}(2/2)/(\mathrm{U}(1) \times \mathrm{U}(1))$ is $\mathrm{SU}(1, 1)/\mathrm{U}(1) \equiv \mathcal{D}^{(1)}$. On the other hand, when one sets the odd variables to zero, the $\mathrm{OSp}(2/2)$ -CS become the $\mathrm{SU}(1, 1)$ -CS, the underlying geometry of which is known to be given by $(\mathcal{D}^{(1)}, \omega_0)$. Furthermore, according to Rothstein's theorem the vector bundle $\mathbb{E}$ is just $\mathbb{F}^*$, the dual of $\mathbb{F}$.

The identification of the remaining data is highly simplified if one makes use of the results of [8]. Indeed, in [8] it has been shown that Rothstein's data of the $\mathrm{OSp}(1/2)$ coadjoint orbit studied there, can be directly read off from the superfunction f generating the supersymplectic superform. More precisely, if one writes f as $f_0 + f_2 + f_4 + \dots$, where f_{2n} designates that component of f which is of order $2n$ in the odd coordinates, then it appears that f_2 assumes the following form:

$$f_2 = -ig_{a\bar{b}}\theta^a\bar{\theta}^b, \quad (4.5)$$

where g is Rothstein's data metric on $\mathbb{E}$; the θ^a 's are the odd supercoordinates of the holomorphic supermanifold considered, which can also be viewed as a (local) frame field of $\mathbb{F}$ over $\mathcal{D}^{(1)}$. The notations in (4.5) are those commonly used in complex geometry, see for example [45].

At this point it is worth anticipating by mentioning that the above considerations are valid only in the *particular complex-analytic* case of a *super Kähler* supermanifold, a notion defined in [8] and rediscussed below. Both $\mathcal{D}^{(1|1)}$ [8] and $\mathcal{D}^{(1|2)}$ are non-trivial examples of such a notion.

We now write the explicit form of g for $\mathcal{D}^{(1|2)}$. If the odd coordinates θ and χ are now

denoted, respectively, by θ^1 and θ^2 , then f_2 in (3.1) is given by:

$$f_2 = i(\tau - b) \frac{\theta^1 \bar{\theta}^1}{(1 - |z|^2)} + i(\tau + b) \frac{\theta^2 \bar{\theta}^2}{(1 - |z|^2)}. \quad (4.6)$$

A simple comparison of (4.5) and (4.6) leads to the following matrix form $||g||$ of g in the frame field of $\mathbb{E}$ over $\mathcal{D}^{(1)}$, which is dual to that of $\mathbb{F}$ given above:

$$||g|| = \begin{pmatrix} \frac{\tau - b}{(1 - |z|^2)} & 0 \\ 0 & \frac{\tau + b}{(1 - |z|^2)} \end{pmatrix}. \quad (4.7)$$

This is a diagonal metric which is clearly Hermitian [45] since we are considering $|b| < \tau$ (typical CS).

It remains now to identify the connection ∇^g compatible with this Hermitian metric. In order to exhibit it, we rederive Rothstein's results (4.1)–(4.2) in our *particular complex-analytic* setting. Once again this task is highly simplified, thanks to the observations of [8] that led to formula (4.5). We recall that Rothstein's formulae (4.1)–(4.3) were derived in the C^∞ case [38].

The complex-analytic counterpart of the even 1-superform α_2 appearing in (4.1), can be obtained from (4.5) as follows:

$$\alpha_2 \equiv -\frac{i}{2}(\delta - \bar{\delta})f_2. \quad (4.8)$$

A direct computation based on (4.5), with g an Hermitian metric on a holomorphic vector bundle $\mathbb{E} \rightarrow M$, leads to:

$$\alpha_2 = \frac{1}{2}g_{a\bar{b}}(\theta^a D\bar{\theta}^b + \bar{\theta}^b D\theta^a), \quad (4.9)$$

where now,

$$D\theta^a = d\theta^a + \Gamma_{ib}^a \theta^b dz^i, \quad D\bar{\theta}^a = d\bar{\theta}^a + \Gamma_{i\bar{b}}^{\bar{a}} \bar{\theta}^{\bar{b}} d\bar{z}^i, \quad (4.10)$$

and

$$\Gamma_{ib}^a = g^{a\bar{c}} \frac{\partial g_{b\bar{c}}}{\partial z^i}. \quad (4.11)$$

One easily recognizes here the Γ_{ib}^a 's as the components of the (canonical) *Hermitian connection* associated to the Hermitian metric g on $\mathbb{E}$ [45]. Notice that these connection components are expressed in the frame field of $\mathbb{E}$, while in the C^∞ case of (4.1)–(4.3) the A_{ib}^a 's are expressed in the frame field of $\mathbb{F}$, dual to that of $\mathbb{E}$. As an endomorphism of a fibre, one is minus the transpose of the other [45]. This explains the sign difference and the change in the notation.

Before stating the final result concerning $(\mathcal{D}^{(1|2)}, \omega)$, let us carry over and evaluate the complex-analytic counterpart of (4.3). From (4.9)–(4.11), a straightforward computation gives,

$$d\alpha_2 = g_{a\bar{b}} D\theta^a D\bar{\theta}^b - g_{a\bar{b}} R_{i\bar{j}c}^a \theta^c \bar{\theta}^b dz^i d\bar{z}^j, \quad (4.12)$$

where

$$R_{jic}^a \equiv \frac{\partial \Gamma_{ic}^a}{\partial \bar{z}^j}. \quad (4.13)$$

From what precedes, we have the following immediate result:

$$\omega_2 = -d\alpha_2, \quad (4.14)$$

where ω_2 is given in (3.5) and $d\alpha_2$ is (4.12) for g the Hermitian metric in (4.7). Hence, in summary, we have proven the following:

Theorem 4.3: *Rothstein's data for $(\mathcal{D}^{(1|2)}, \omega)$ are: $(\mathcal{D}^{(1)}, \omega_0, \mathbb{E}, g, \nabla^g)$ where $\mathcal{D}^{(1)}$ is the unit disc, ω_0 is the $SU(1, 1)$ -invariant two-form on $\mathcal{D}^{(1)}$ (3.4), $\mathbb{E} = \mathbb{F}^*$ is a rank 2 trivial holomorphic vector bundle over $\mathcal{D}^{(1)}$, g is the Hermitian metric (4.7) on $\mathbb{E}$ and ∇^g is the corresponding (canonical) Hermitian connection (the components of which can be explicitly evaluated using (4.11)).*

Notice now that the symplectic $(\mathcal{D}^{(1)}, \omega_0)$ is moreover a Kähler manifold, since $\omega_0 = -i\partial\bar{\partial}f_0$, where f_0 , the odd-coordinates-independent part of f in (3.1), is a Kähler potential for ω_0 ; $\partial = dz \frac{\partial}{\partial z}$. Hence, one clearly sees that $(\mathcal{D}^{(1|2)}, \omega)$ is a non-trivial example of the following definition of a super Kähler supermanifold:

Definition 4.4: *A super Kähler supermanifold $(M, \mathcal{A}_M, \omega)$ is a holomorphic supersymplectic supermanifold, whose Rothstein's data, $(M, \omega_0, \mathbb{E}, g, \nabla^g)$, are such that (M, ω_0) is a Kähler manifold, $(\mathbb{E}, g)$ is a holomorphic Hermitian vector bundle over M and ∇^g is the canonical Hermitian connection.*

As already mentioned the present situation allows us to go beyond Rothstein's theorem [38]. Indeed, ω_4 in (3.3)–(3.6) can also be rewritten in terms of Rothstein's data of $(\mathcal{D}^{(1|2)}, \omega)$ obtained above. This is achieved by simply noticing that in (3.1):

$$f_4 = \frac{1}{4\tau}(f_2)^2. \quad (4.15)$$

Then,

$$\alpha_4 \equiv -\frac{i}{2}(\delta - \bar{\delta})f_4 = \frac{1}{2\tau}f_2\alpha_2, \quad (4.16)$$

such that,

$$\omega_4 \equiv -d\alpha_4 = \frac{1}{2\tau} [f_2\omega_2 - df_2\alpha_2]. \quad (4.17)$$

A simple computation based on (4.5), (4.9) and (4.12), leads to the following:

$$\omega_4 = \frac{i}{2\tau} g_{a\bar{b}} g_{c\bar{d}} [\theta^a \bar{\theta}^d D\theta^c D\bar{\theta}^b + \theta^a \bar{\theta}^b (D\theta^c D\bar{\theta}^d - R_{i\bar{j}e}^c \theta^e \bar{\theta}^d dz^i d\bar{z}^j)]. \quad (4.18)$$

When g is the Hermitian metric in (4.7), ω_4 above is exactly the ω_4 of (3.6). The present extension of Rothstein's theorem is mainly based on the observation made in (4.15). This is in fact an intrinsic property of super Kähler coadjoint orbits of some simple Lie supergroups. For such orbits, Rothstein's data determine the complete super Kähler form. A detailed description of a general framework will be given elsewhere.

In analogy with the non-super case, and in view of (3.3) it is worth calling the real superfunction f in (3.1) a super Kähler potential for the super Kähler form ω given in (3.3)–(3.6). Such a potential is defined up to the addition of a superholomorphic or/and an anti-superholomorphic function on $\mathcal{D}^{(1|2)}$. Clearly, Rothstein's data for a super Kähler supermanifold are encoded in its super Kähler potential $f = f_0 + f_2 + \dots$. Indeed, the body f_0 of f is a Kähler potential for the Kähler manifold (M, ω_0) , while f_2 provides the Hermitian structure on the holomorphic bundle $\mathbb{E}$ (see 4.5).

The description of super Kähler geometry given above is sufficient for our purpose. However, it is worth mentioning that Definition 4.4 can be made much more precise. Indeed, a deeper analysis of Rothstein's data for $(\mathcal{D}^{(1|2)}, \omega)$ shows that they determine an Einstein-Hermitian vector bundle [45] (see Section 7, 7.3). This very interesting observation deserves further investigations. More details will be given in a forthcoming publication.

The next section addresses the geometric quantization of $(\mathcal{D}^{(1|2)}, \omega)$. The super Kähler character of the latter leads naturally to the existence of a super Kähler polarization, which makes the complete quantization program successful.

5 Geometric quantization

By *geometric quantization* we refer here to the celebrated method, independently devised by Kostant [17] and Souriau [18], which associates to a given classical mechanics a quantum counterpart. Kostant-Souriau quantization procedure meets Kirillov's *method of orbits* [16] when the classical mechanics is described by a coadjoint orbit of a Lie group G . The quantum output comes then in the form of a Unitary Irreducible Representation

(UIR) of the considered group G . This is the case we are interested in super extending. From now on we will focus on this kind of situations.

In practice, geometric quantization proceeds in two steps. The first one, called *pre-quantization*, consists in exhibiting a complex line bundle over $(M \equiv G/H, \omega)$, with Hermitian structure and compatible connection ∇ , such that $\text{curv}\nabla = \omega$. Such a line bundle exists whenever $[\omega]$ is an integral cohomology class (integrality condition). When lifted to this line bundle, the transitive (and symplectic) action of G on M gives rise to a unitary but reducible representation of G . The second step consists in using a *polarization* in order to select an irreducible subrepresentation of the prequantum representation. More precisely, the group action is restricted to the subspace of those L^2 sections of the prequantum line bundle which are covariantly constant along the vector fields generating the polarization. More details concerning the general procedure of geometric quantization can be found in [46, 47].

Super-prequantization was partly developed by Kostant [28]. Tuynman [48] completed that construction by equipping Kostant's super-prequantum bundle with a super Hermitian structure compatible with the connection. We will follow here that construction, assuming that the reader is at least familiar with Kostant's work. On the other hand, because of the lack of a notion of polarization, the second part of the program was not considered in [28]. Here, as a super Kähler supermanifold $(\mathcal{D}^{(1|2)}, \omega)$ is naturally equipped with a *super Kähler polarization* that allows us to carry out the whole quantization program.

Before giving the details of the construction, it is worth mentioning that the notion of a polarization in the super context and in connection with geometric quantization already appeared in the literature. For instance, in the real case, a general definition of a polarization was given in [48]; it was then used to quantize the BRST charge. On the other hand, a super Kähler polarization was introduced in [49] in order to quantize a field theory with fermionic degrees of freedom, i.e. an infinite dimensional flat phase space. Here we use a general notion in the complex-analytic case which allows quantizing non-trivial super phase spaces, such as coadjoint orbits.

Finally, let us mention that recently another quantization method, namely the deformation quantization, has been extended to the super context in [50]; this method was applied in particular to the super unit disc $\mathcal{D}^{(1|1)}$. Geometric quantization of the latter is considered in [9].

5.1 Super prequantization

Following Kostant's general scheme [28], in order to prequantize $(\mathcal{D}^{(1|2)}, \omega)$, one needs to exhibit a complex *line bundle sheaf* $\mathcal{L}^{(1|2)}$ with connection ∇ over $\mathcal{D}^{(1|2)}$, such that $\text{curv}\nabla = \omega$. Such a line bundle sheaf exists if and only if there exists a complex line bundle $\mathbb{L}$ with connection ∇_0 over $\mathcal{D}^{(1)}$, such that $\text{curv}\nabla_0 = \omega_0$ (i.e. *iff* $(\mathcal{D}^{(1)}, \omega_0)$ is prequantizable) [28]. Since our ω_0 is exact, it is known that such an $\mathbb{L}$ always exists. It is then not hard to see that,

$$\mathcal{L}^{(1|2)} \equiv \mathcal{L}^{(1)} \otimes \mathcal{A}_{\mathbb{C}}^{(1|2)} \cong \mathcal{A}_{\mathbb{C}}^{(1|2)}, \quad (5.1)$$

where $\mathcal{L}^{(1)}$ is the sheaf of C^∞ sections of $\mathbb{L}$.

To the classical observables in (3.9) one can associate the so-called prequantum operators, which act in the space of C^∞ sections of $\mathcal{L}^{(1|2)}$. These operators are obtained using the following formula [28],

$$\hat{H} \equiv -i\nabla_{X_H} + H, \quad \text{where} \quad \nabla_{X_H} = X_H - iX_H \lrcorner \vartheta, \quad (5.2)$$

and

$$\begin{aligned} \vartheta \equiv -i\delta f = & -2i\tau \frac{\bar{z}dz}{1-|z|^2} \left[1 + i\frac{\tau-b}{2\tau} \frac{\bar{\theta}\theta}{1-|z|^2} + i\frac{\tau+b}{2\tau} \frac{\bar{\chi}\chi}{1-|z|^2} - \frac{\tau^2-b^2}{2\tau^2} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1-|z|^2)^2} \right] \\ & + (\tau-b) \frac{d\theta}{1-|z|^2} \frac{\bar{\theta}}{1-|z|^2} \left[1 + i\frac{\tau+b}{2\tau} \frac{\bar{\chi}\chi}{1-|z|^2} \right] + (\tau+b) \frac{d\chi}{1-|z|^2} \frac{\bar{\chi}}{1-|z|^2} \left[1 + i\frac{\tau-b}{2\tau} \frac{\bar{\theta}\theta}{1-|z|^2} \right] \end{aligned} \quad (5.3)$$

is a 1-superform potential for the connection ∇ equipping the prequantum bundle $\mathcal{L}^{(1|2)}$; it is such that $\omega = -d\vartheta$. A straightforward computation gives,

$$\begin{aligned} \hat{B} &= \frac{\theta}{2} \frac{\partial}{\partial \theta} - \frac{\bar{\theta}}{2} \frac{\partial}{\partial \bar{\theta}} - \frac{\chi}{2} \frac{\partial}{\partial \chi} + \frac{\bar{\chi}}{2} \frac{\partial}{\partial \bar{\chi}} + b, \\ \hat{K}_0 &= z \frac{\partial}{\partial z} - \bar{z} \frac{\partial}{\partial \bar{z}} + \frac{\theta}{2} \frac{\partial}{\partial \theta} - \frac{\bar{\theta}}{2} \frac{\partial}{\partial \bar{\theta}} + \frac{\chi}{2} \frac{\partial}{\partial \chi} - \frac{\bar{\chi}}{2} \frac{\partial}{\partial \bar{\chi}} + \tau, \\ \hat{K}_+ &= z^2 \frac{\partial}{\partial z} - \frac{\partial}{\partial \bar{z}} + z\theta \frac{\partial}{\partial \theta} + z\chi \frac{\partial}{\partial \chi} + 2\tau z, \\ \hat{K}_- &= \frac{\partial}{\partial z} - \bar{z}^2 \frac{\partial}{\partial \bar{z}} - \bar{z}\bar{\theta} \frac{\partial}{\partial \bar{\theta}} - \bar{z}\bar{\chi} \frac{\partial}{\partial \bar{\chi}}, \\ \hat{V}_+ &= -i \left(\frac{\tau-b}{2\tau} \right) z\theta \frac{\partial}{\partial z} - \left(\frac{\tau+b}{2\tau} \right) \bar{\chi} \frac{\partial}{\partial \bar{z}} + \frac{\partial}{\partial \bar{\theta}} + i \left(z + \left(\frac{\tau-b}{2\tau} \right) \chi\theta \right) \frac{\partial}{\partial \chi} - i(\tau-b)\theta, \\ \hat{W}_- &= -i \left(\frac{\tau+b}{2\tau} \right) \chi \frac{\partial}{\partial z} - \left(\frac{\tau-b}{2\tau} \right) \bar{z}\bar{\theta} \frac{\partial}{\partial \bar{z}} + i \frac{\partial}{\partial \theta} + \left(\bar{z} - \left(\frac{\tau-b}{2\tau} \right) \bar{\theta}\bar{\chi} \right) \frac{\partial}{\partial \bar{\chi}}, \end{aligned} \quad (5.4)$$

$$\begin{aligned}\hat{V}_- &= -i \left(\frac{\tau-b}{2\tau} \right) \theta \frac{\partial}{\partial z} - \left(\frac{\tau+b}{2\tau} \right) \bar{z} \bar{\chi} \frac{\partial}{\partial \bar{z}} + \left(\bar{z} + \left(\frac{\tau+b}{2\tau} \right) \bar{\theta} \bar{\chi} \right) \frac{\partial}{\partial \bar{\theta}} + i \frac{\partial}{\partial \chi}, \\ \hat{W}_+ &= -i \left(\frac{\tau+b}{2\tau} \right) z \chi \frac{\partial}{\partial z} - \left(\frac{\tau-b}{2\tau} \right) \bar{\theta} \frac{\partial}{\partial \bar{z}} + i \left(z - \left(\frac{\tau+b}{2\tau} \right) \chi \theta \right) \frac{\partial}{\partial \theta} + \frac{\partial}{\partial \bar{\chi}} - i(\tau+b)\chi.\end{aligned}$$

One easily verifies that these operators close to the $\mathfrak{osp}(2/2)$ Lie superalgebra. They provide thus a representation of $\mathfrak{osp}(2/2)$ in the space of C^∞ sections of $\mathcal{L}^{(1|2)}$.

In the next subsection, using a natural *invariant super Kähler polarization* on $\mathcal{D}^{(1|2)}$ we will select a subsheaf $\mathcal{L}_p^{(1|2)}$ of $\mathcal{L}^{(1|2)}$; $\mathcal{L}_p^{(1|2)}$ will then be equipped with a ∇ -compatible *super Hermitian structure*. When restricted to $\mathcal{L}_p^{(1|2)}$ the above prequantum representation will reduce to a *super unitary irreducible representation* of $\mathfrak{osp}(2/2)$.

5.2 Super polarization

As for the Kähler unit disc $\mathcal{D}^{(1)}$, a natural super polarization, called here *super Kähler* and denoted $\mathcal{P}$, exists on $\mathcal{D}^{(1|2)}$. It is spanned by the vector superfields $\frac{\partial}{\partial z}, \frac{\partial}{\partial \theta}$ and $\frac{\partial}{\partial \chi}$. One easily verifies that this is actually a good candidate for a super polarization. Indeed, $\mathcal{P}$ fulfills all the required conditions, namely,

- (i) $\mathcal{P}$ is involutive;
- (ii) $\mathcal{P}$ is maximal isotropic, i.e. $\omega(Z, Y) = 0, \forall Z, Y \in \mathcal{P}$ and $\dim \mathcal{P} = (1|2) = \dim \mathcal{D}^{(1|2)}$ as a holomorphic supermanifold.

Moreover, one can easily verify that,

- (iii) $\mathcal{P} \cap \bar{\mathcal{P}} = \{0\}$;
- (iv) $\mathcal{P}$ is invariant, namely, for X_H one of the Hamiltonian vector fields in (3.12), $[X_H, Z] \in \mathcal{P}, \forall Z \in \mathcal{P}$ (brackets denote here a commutator or an anticommutator).

Property (iii) confirms, in agreement with the results of Section 4, that $\mathcal{P}$ deserves to be called a *super Kähler polarization* for $\mathcal{D}^{(1|2)}$. On the other hand, property (iv) means that the Hamiltonian flows of the classical observables (3.9) preserve $\mathcal{P}$; the importance of this property will be stressed soon. Moreover, as it will be shown at the end of this section, $\mathcal{P}$ is *positive*. This property ensures that the final representation space is non-trivial. In summary, $\mathcal{P}$ is a positive invariant super Kähler polarization.

This polarization is now used in order to select a subsheaf $\mathcal{L}_p^{(1|2)}$ of $\mathcal{L}^{(1|2)}$. The latter consists of the sections $\psi(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi})$ of $\mathcal{L}^{(1|2)}(\mathcal{D}^{(1)})$ which are covariantly constant along $\overline{\mathcal{P}}$, i.e. those sections ψ which are such that:

$$\nabla_{\frac{\partial}{\partial \bar{z}}} \psi = 0, \quad \nabla_{\frac{\partial}{\partial \bar{\theta}}} \psi = 0, \quad \text{and} \quad \nabla_{\frac{\partial}{\partial \bar{\chi}}} \psi = 0, \quad (5.5)$$

where, from now on, the covariant superderivation is taken as in (5.2) but with ϑ , the superpolarization-adapted 1-superform potential for ∇ , replaced by the *real* 1-superform potential α given by,

$$\begin{aligned} \alpha &\equiv -\frac{i}{2}(\delta - \bar{\delta})f \\ &= i\tau \frac{z d\bar{z} - \bar{z} dz}{1 - |z|^2} \left[1 + i\frac{\tau - b}{2\tau} \frac{\bar{\theta}\theta}{1 - |z|^2} + i\frac{\tau + b}{2\tau} \frac{\bar{\chi}\chi}{1 - |z|^2} - \frac{\tau^2 - b^2}{2\tau^2} \frac{\bar{\theta}\bar{\chi}\chi\theta}{(1 - |z|^2)^2} \right] \\ &\quad + \frac{\tau - b}{2} \frac{d\theta\bar{\theta} + d\bar{\theta}\theta}{1 - |z|^2} \left[1 + i\frac{\tau + b}{2\tau} \frac{\bar{\chi}\chi}{1 - |z|^2} \right] + \frac{\tau + b}{2} \frac{d\chi\bar{\chi} + d\bar{\chi}\chi}{1 - |z|^2} \left[1 + i\frac{\tau - b}{2\tau} \frac{\bar{\theta}\theta}{1 - |z|^2} \right]. \end{aligned} \quad (5.6)$$

In (5.6) $f = f(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi})$ is the super Kähler potential (3.1). Solutions to (5.5) are of the form

$$\psi(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi}) = \exp(-f/2) \phi(z, \theta, \chi), \quad (5.7)$$

where ϕ is a superholomorphic section of $\mathcal{L}^{(1|2)}$, and $\exp(-f/2) \equiv \mathcal{N}'$ (see (2.28) and (3.1)). Notice from (5.7) that $\mathcal{L}_p^{(1|2)}$ is isomorphic to $\mathcal{A}^{(1|2)}$, the superstructure sheaf of $\mathcal{D}^{(1|2)}$ viewed as a holomorphic supermanifold. Moreover, property **(iv)** above ensures that the action of the prequantum operators (5.4) in $\mathcal{L}_p^{(1|2)}$ leaves the latter invariant.

Let us now equip $\mathcal{L}_p^{(1|2)}(\mathcal{D}^{(1)})$ with a super Hermitian structure $(\cdot, \cdot)$. For $\psi = \psi_{\bar{0}} + \psi_{\bar{1}}$ and $\psi' = \psi'_0 + \psi'_1 \in \mathcal{L}_p^{(1|2)}(\mathcal{D}^{(1)}) = \mathcal{L}_p^{(1|2)}(\mathcal{D}^{(1)})_{\bar{0}} \oplus \mathcal{L}_p^{(1|2)}(\mathcal{D}^{(1)})_{\bar{1}}$, this is given by:

$$(\psi', \psi) \equiv \overline{\psi'} \psi = \overline{\psi'_0} \psi_{\bar{0}} + \overline{\psi'_1} \psi_{\bar{1}} + \overline{\psi'_0} \psi_{\bar{1}} + \overline{\psi'_1} \psi_{\bar{0}}. \quad (5.8)$$

The latter clearly satisfies (2.13). Notice however that $(\cdot, \cdot)$ is *not* homogeneous, i.e. it is not of the form (2.14). Moreover, it is not hard to verify that the super connection ∇ on $\mathcal{L}^{(1|2)}$ is compatible with (5.8), i.e.

$$X(\psi', \psi) = (\nabla_X \psi', \psi) + (-1)^{\epsilon(\psi')\epsilon(X)} (\psi', \nabla_X \psi), \quad (5.9)$$

where ψ' is now a homogeneous section of $\mathcal{L}_p^{(1|2)}(\mathcal{D}^{(1)})$ of parity $\epsilon(\psi')$, X is a real homogeneous vector superfield on $\mathcal{D}^{(1|2)}$ of parity $\epsilon(X)$, and $\nabla_X = X - iX \lrcorner \alpha$, for α given in (5.6).

As for usual geometric quantization in presence of a Kähler polarization, using the super Hermitian structure (5.8), the superspace of sections ψ of $\mathcal{L}_p^{(1|2)}(\mathcal{D}^{(1)})$ can be equipped with a super-inner product, given by:

$$\langle\langle \psi', \psi \rangle\rangle_{\tau, b} \equiv \int_{\mathcal{D}^{(1|2)}} (\psi', \psi) d\mu, \quad (5.10)$$

where $d\mu = d\mu(z, \bar{z}, \theta, \bar{\theta}, \chi, \bar{\chi})$ is the $\text{OSp}(2/2)$ -invariant super measure on $\mathcal{D}^{(1|2)}$ obtained from ω in (3.19). Because of the isomorphism mentioned after (5.7), this super-inner product can be understood as an inner product on the space of sections ϕ of $\mathcal{A}^{(1|2)}$ (i.e. the space of superholomorphic sections of $\mathcal{L}^{(1|2)}$). Hence, using (5.7), we can write

$$\langle\langle \phi', \phi \rangle\rangle_{\tau, b} = \int_{\mathcal{D}^{(1|2)}} e^{-f} (\phi', \phi) d\mu. \quad (5.11)$$

Let us now investigate the status of this super-inner product from the Hilbert space point of view.

Since $|b| < \tau$, one can always write a superholomorphic function $\phi(z, \theta, \chi)$ on $\mathcal{D}^{(1|2)}$ as follows:

$$\phi(z, \theta, \chi) = \phi_1(z) + \theta \sqrt{\tau - b} \phi_2(z) + \chi \sqrt{\tau + b} \phi_3(z) + \chi \theta \sqrt{\frac{(\tau^2 - b^2)(2\tau + 1)}{2\tau}} \phi_4(z), \quad (5.12)$$

where $\phi_i, i = 1, \dots, 4$, are holomorphic functions on $\mathcal{D}^{(1)}$.

The super integration in (5.11) can be partially carried out. Indeed, replacing (3.1), (3.19), (5.8) and (5.12) in (5.11), and using Berezin integration over the odd Grassmann variables [22], namely, the only non-zero integral being $\int d\theta d\bar{\theta} d\chi d\bar{\chi} (\bar{\chi} \chi \bar{\theta} \theta) = 1$, one obtains:

$$\langle\langle \phi', \phi \rangle\rangle_{\tau, b} = \langle \phi'_1, \phi_1 \rangle_{k=\tau} + i \langle \phi'_2, \phi_2 \rangle_{k=\tau+\frac{1}{2}} + i \langle \phi'_3, \phi_3 \rangle_{k=\tau+\frac{1}{2}} + \langle \phi'_4, \phi_4 \rangle_{k=\tau+1} \in \mathbb{C}, \quad (5.13)$$

where

$$\langle \phi', \phi \rangle_k \equiv \frac{2k-1}{\pi} \int_{\mathcal{D}^{(1)}} \overline{\phi'(z)} \phi(z) \frac{dz d\bar{z}}{(1-|z|^2)^{2-2k}}, \quad \text{for } k > \frac{1}{2}, \quad (5.14)$$

is the usual inner product on the representation space of the holomorphic (positive) discrete series $D(k)$ of $\text{SU}(1, 1)$, which arise through geometric quantization of the unit disc $(\mathcal{D}^{(1)}, \omega_0)$ [12, 51, 52].

We can now define a natural notion of square integrability of superholomorphic sections of a prequantum bundle sheaf, and thus that of a *super Hilbert space*. First observe that

$\langle\langle \cdot, \cdot \rangle\rangle_{\tau, b}$ is an *even super Hermitian form* on the space of superholomorphic sections of $\mathcal{L}^{(1|2)}$. Indeed,

$$\overline{\langle\langle \phi', \phi \rangle\rangle_{\tau, b}} = (-1)^{\epsilon(\phi')\epsilon(\phi)} \langle\langle \phi, \phi' \rangle\rangle_{\tau, b}, \quad (5.15)$$

and moreover,

$$\langle\langle \phi', \phi \rangle\rangle_{\tau, b} = \langle\langle \phi'_0, \phi_0 \rangle\rangle_{\tau, b, \bar{0}} + i \langle\langle \phi'_1, \phi_1 \rangle\rangle_{\tau, b, \bar{1}}. \quad (5.16)$$

The new quantities and notation in the above equation can be easily identified in terms of the old ones appearing in (5.7), (5.13) and (5.14). A simple comparison of (5.15) and (5.16) with respectively (2.13) and (2.14) confirms our claim. This suggests the following natural definition:

Definition 5.1: A superholomorphic section $\phi = \phi_0 + \phi_1$ of $\mathcal{L}^{(1|2)}$ is said to be square integrable if both ϕ_0 and ϕ_1 are respectively square integrable with respect to the Hermitian forms $\langle\langle \cdot, \cdot \rangle\rangle_{\tau, b, \bar{0}}$ and $\langle\langle \cdot, \cdot \rangle\rangle_{\tau, b, \bar{1}}$.

A definition of a *super Hilbert space*, different from those proposed in [27, 53, 54], immediately follows.

Definition 5.2: A super Hilbert space is a pair $(\mathcal{H}, \langle\langle \cdot, \cdot \rangle\rangle)$, where $\mathcal{H} \equiv \mathcal{H}_0 \oplus \mathcal{H}_1$ is a superspace equipped with a super Hermitian form, $\langle\langle \cdot, \cdot \rangle\rangle = \langle\langle \cdot, \cdot \rangle\rangle_0 + i \langle\langle \cdot, \cdot \rangle\rangle_1$, such that $(\mathcal{H}_0, \langle\langle \cdot, \cdot \rangle\rangle_0)$ and $(\mathcal{H}_1, \langle\langle \cdot, \cdot \rangle\rangle_1)$ are both Hilbert spaces.

According to these definitions, the space of superholomorphic sections of $\mathcal{L}^{(1|2)}$ equipped with the super Hermitian form (5.13) can be turned into a super Hilbert space simply by taking $\phi_i, i = 1, \dots, 4$, to be L^2 functions on the unit disc with respect to the corresponding Hermitian forms in (5.12), namely, that $\|\phi_1\|_{k=\tau}^2 < \infty$, $\|\phi_2\|_{k=\tau+\frac{1}{2}}^2 < \infty$, $\|\phi_3\|_{k=\tau+\frac{1}{2}}^2 < \infty$ and $\|\phi_4\|_{k=\tau+1}^2 < \infty$. Hence, in these conditions, $\phi(z, \theta, \chi)$ in (5.12) can be called an L^2 superholomorphic section of $\mathcal{L}^{(1|2)}$ with respect to the super Hermitian form of (5.11). The obtained super Hilbert space, denoted $\mathcal{H}_{\tau, b}$, constitutes then the representation superspace carrying an *explicit* realization of the typical irreducible representation $V(\tau, b)$ of $\mathfrak{osp}(2/2)$.

Remark 5.3: We could have equipped $\mathcal{L}^{(1|2)}(\mathcal{D}^{(1)})$ with an *even* super Hermitian structure $((\cdot, \cdot))$ instead of the one of indefinite parity introduced in (5.8). The former would assume the following form $((\psi', \psi)) \equiv \overline{\psi'_0} \psi_0 + \overline{\psi'_1} \psi_1$. In this case (5.9) will be true only modulo *odd* quantities. However, $\langle\langle \cdot, \cdot \rangle\rangle_{\tau, b} \equiv \int_{\mathcal{D}^{(1|2)}} (\cdot, \cdot) d\mu =$

$\int_{\mathcal{D}^{(1|2)}} ((\cdot, \cdot)) d\mu$, since $((\cdot, \cdot))$ and $(\cdot, \cdot)$ differ only by *odd* quantities which disappear when one integrates over anti-commuting variables using Berezin integration. For the same reason, all the results following (5.10) will still hold true.

Remark 5.4: The important result in (5.13)–(5.14) follows from the special form adopted for ϕ in (5.12), and the choice of normalization made for $d\mu$ in Section 3.3, namely, $\int_{\mathcal{D}^{(1|2)}} \exp(-f) d\mu = 1$.

The generators of $\mathfrak{osp}(2/2)$ are represented in $\mathcal{H}_{\tau,b}$ by the super holomorphic restrictions of the prequantum operators (5.4). More precisely,

$$\begin{aligned}
\widehat{B} &= \frac{\theta}{2} \frac{\partial}{\partial \theta} - \frac{\chi}{2} \frac{\partial}{\partial \chi} + b, \\
\widehat{K}_0 &= z \frac{\partial}{\partial z} + \frac{\theta}{2} \frac{\partial}{\partial \theta} + \frac{\chi}{2} \frac{\partial}{\partial \chi} + \tau, \\
\widehat{K}_+ &= z^2 \frac{\partial}{\partial z} + z\theta \frac{\partial}{\partial \theta} + z\chi \frac{\partial}{\partial \chi} + 2\tau z, \\
\widehat{K}_- &= \frac{\partial}{\partial z}, \\
\widehat{V}_+ &= -i \left(\frac{\tau - b}{2\tau} \right) z\theta \frac{\partial}{\partial z} + i \left(z + \left(\frac{\tau - b}{2\tau} \right) \chi\theta \right) \frac{\partial}{\partial \chi} - i(\tau - b)\theta, \\
\widehat{W}_- &= -i \left(\frac{\tau + b}{2\tau} \right) \chi \frac{\partial}{\partial z} + i \frac{\partial}{\partial \theta}, \\
\widehat{V}_- &= -i \left(\frac{\tau - b}{2\tau} \right) \theta \frac{\partial}{\partial z} + i \frac{\partial}{\partial \chi}, \\
\widehat{W}_+ &= -i \left(\frac{\tau + b}{2\tau} \right) z\chi \frac{\partial}{\partial z} + i \left(z - \left(\frac{\tau + b}{2\tau} \right) \chi\theta \right) \frac{\partial}{\partial \theta} - i(\tau + b)\chi.
\end{aligned} \tag{5.17}$$

The obtained representation is *super unitary*. Indeed, in agreement with (5.9), the *superadjoint* (or *super Hermitian conjugate*) $\widehat{X}^\dagger$ of an operator $\widehat{X}$ acting in $\mathcal{H}_{\tau,b}$ is defined as follows:

$$\langle\langle \widehat{X}^\dagger \phi', \phi \rangle\rangle_{\tau,b} = (-1)^{\epsilon(\phi')\epsilon(X)} \langle\langle \phi', \widehat{X} \phi \rangle\rangle_{\tau,b}, \tag{5.18}$$

for ϕ' a homogeneous superholomorphic section of $\mathcal{L}^{(1|2)}$ and $\widehat{X}$ a homogeneous operator. Hence, the quantum counterpart $\widehat{H}$ of a *real* classical observable H^{cl} is a *self superadjoint* operator acting in $\mathcal{H}_{\tau,b}$, i.e. $\widehat{H}^\dagger = \widehat{H}$. Specifically, since both B^{cl} and K_0^{cl} are real (see (3.10)), the associated quantum operators in (5.17) are self superadjoint:

$$\widehat{B}^\dagger = \widehat{B} \quad \text{and} \quad \widehat{K}_0^\dagger = \widehat{K}_0. \tag{5.19}$$

On the other hand, the reality of $(K_+^{cl} + K_-^{cl})$, $(V_+^{cl} + iW_-^{cl})$ and of similar combinations that can be obtained from (3.10), leads to the following,

$$(\widehat{K}_+)^{\dagger} = \widehat{K}_-, \quad (\widehat{V}_+)^{\dagger} = i\widehat{W}_-, \quad \text{and} \quad (\widehat{V}_-)^{\dagger} = i\widehat{W}_+. \quad (5.20)$$

As it should be, these results are in perfect agreement with the relations we started with at the level of the abstract representation theory (compare (5.18)–(5.20), with (B.10), (B.13) and (B.14)).

Geometric quantization of $(\mathcal{D}^{(1|2)}, \omega)$ is hence completed. It remains now to integrate the obtained super unitary irreducible representation of $\mathfrak{osp}(2/2)$ to a representation of $\mathrm{OSp}(2/2)$ (see Section 7, 7.5). The integrability condition, which we assumed from the beginning, ensures that such a procedure leads actually to a non-trivial representation of $\mathrm{OSp}(2/2)$. Indeed, recall that we started with an *integrable* typical $\mathfrak{osp}(2/2)$ -module $V(\tau, b)$, i.e. both b and $\tau \in \frac{1}{2}\mathbb{N}$. Moreover, because of (5.14), we consider $\tau > \frac{1}{2}$. This last condition is a direct consequence of the so-called metaplectic correction to geometric quantization [46], which we do not consider here.

Finally we briefly discuss the positivity of our super Kähler polarization $\mathcal{P}$. In the non-super context, the positivity of a Kähler polarization ensures that the unitary irreducible representation obtained is non-trivial, i.e. the corresponding Hilbert space does not reduce to the zero function [46]. If a super extension of the positivity exists, then $\mathcal{P}$ is positive, since $\mathcal{H}_{\tau, b}$ is clearly non-trivial. An analysis of the present situation and of the one considered in [9] lead to a natural supergeometric definition of the positivity of a super Kähler polarization. This will be given elsewhere.

6 Atypical $\mathrm{OSp}(2/2)$ -CS and associated orbits

We now briefly discuss the main results concerning the $\mathrm{OSp}(2/2)$ atypical representations: the atypical coherent states and their underlying supergeometry. Recall that the atypical representations occur when $|b| = \tau$. It is then not hard to see that by taking $b \rightarrow \tau$ or $b \rightarrow -\tau$, almost all the results obtained in the previous sections reduce to atypical analogs. Here we emphasize the main features of the case $b = -\tau$ (one obtains exactly the same results when $b = \tau$).

If $b = -\tau$, the lowest weight vector is $|- \tau, \tau, \tau\rangle \equiv |\tau, \tau\rangle$. As it is shown in Appendix B, $W_+|\tau, \tau\rangle$ is a primitive vector generating an $\mathfrak{osp}(2/2)$ -submodule of zero-norm states.

Hence, $|\tau, \tau\rangle$ satisfies not only (2.5)–(2.6), but it is also such that

$$W_+|\tau, \tau\rangle = 0. \quad (6.1)$$

The appropriate irreducible $\mathfrak{osp}(2/2)$ -module is no longer $V(\tau, b)$, given in (2.9), but the one given in (2.11). Instead of having four families of states, as in (2.10) or (B.1), we have only two:

$$K_+^m|\tau, \tau\rangle \quad \text{and} \quad K_+^m V_+^m|\tau, \tau\rangle, \quad m \geq 0. \quad (6.2)$$

In other words,

$$U(\tau) \equiv V(\tau, -\tau)/V'(\tau, -\tau) \equiv \text{span}\{|\tau, \tau + m\rangle, |\tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle, m \in \mathbb{N}\}. \quad (6.3)$$

The same techniques as in Section 2 lead to the following atypical CS in $\tilde{U}(\tau)$. Starting with $|\tau, \tau\rangle$, these CS appear to be parametrized only by two variables. Indeed, as a consequence of (6.1),

$$\begin{aligned} |a, \theta, \chi\rangle &\equiv \mathcal{M} \exp(aK_+ + \theta V_+ + \chi W_+)|\tau, \tau\rangle, \\ &\equiv \mathcal{M} \exp(zK_+ + \theta V_+)|\tau, \tau\rangle \equiv |z, \theta\rangle, \end{aligned} \quad (6.4)$$

where $a \in \mathcal{B}_0$, and $\theta, \chi \in \mathcal{B}_1$ such that $z = a - \frac{1}{2\tau}\chi\theta \in \mathbb{C}$. A simple comparison of (6.4) with the corresponding equation for the typical CS in Section 2, shows that the atypical CS are simply obtained from the typical ones by taking $b = -\tau$; clearly they do not depend on χ . More precisely,

$$\begin{aligned} |z, \theta\rangle &= \mathcal{M} \left[\sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m)}{m!\Gamma(2\tau)}} z^m |\tau, \tau + m\rangle \right. \\ &\quad \left. + \theta \sqrt{2\tau} \sum_{m=0}^{\infty} \sqrt{\frac{\Gamma(2\tau+m+1)}{m!\Gamma(2\tau+1)}} z^m |\tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle \right], \end{aligned} \quad (6.5)$$

where the real normalization constant is

$$\mathcal{M} = (1 - |z|^2)^\tau \left[1 + i\tau \frac{\bar{\theta}\theta}{1 - |z|^2} \right]. \quad (6.6)$$

At this point it is worth noticing, in connection with Remark 2.7, that the atypical CS are nothing but $\text{OSp}(1/2)$ -CS [34, 8], where here $\text{OSp}(1/2)$ is the subsupergroup of $\text{OSp}(2/2)$ generated by $\{K_0, K_\pm, F_\pm \equiv (V_\pm + W_\pm)/\sqrt{2}\}$. More precisely,

$$|z, \theta\rangle = \mathcal{M} \exp(zK_+ + \sqrt{2}\theta F_+)|\tau, \tau\rangle; \quad (6.7)$$

once again this is a direct consequence of (6.1). Hence, up to the above rescaling of θ , the analysis of the supergeometry underlying the $\text{OSp}(2/2)$ atypical CS, and its geometric quantization reduce *almost* to those already studied in [8, 9]. The main differences are discussed below.

The variables z and θ above parametrize the $N = 1$ super unit disc $\text{OSp}(1/2)/\text{U}(1) \equiv \mathcal{D}^{(1|1)}$ [8]. The super Kähler potential on $\mathcal{D}^{(1|1)}$ is given by,

$$f(z, \bar{z}, \theta, \bar{\theta}) = \log |\langle \tau, \tau | \bar{z}, \bar{\theta} \rangle|^{-2} = -2\tau \left[\log(1 - |z|^2) - i \frac{\bar{\theta}\theta}{1 - |z|^2} \right], \quad (6.8)$$

and the super Kähler superform is simply,

$$\omega = \frac{-2i\tau}{(1 - |z|^2)^2} \left[1 + i\bar{\theta}\theta \frac{1 + |z|^2}{(1 - |z|^2)} \right] dz d\bar{z} - \frac{2\tau}{1 - |z|^2} d\theta d\bar{\theta} - \frac{2\tau}{(1 - |z|^2)^2} [\theta \bar{z} dz d\bar{\theta} - \bar{\theta} z d\theta d\bar{z}]. \quad (6.9)$$

The $\text{osp}(2/2)$ classical observables are obtained from (3.9) by taking $b = -\tau$, or equivalently by formally setting $\chi = 0$. When evaluated as in the typical case, the Hamiltonian vector superfields assume now the following form,

$$X_B = i \frac{\theta}{2} \frac{\partial}{\partial \theta} - i \frac{\bar{\theta}}{2} \frac{\partial}{\partial \bar{\theta}}, \quad X_{K_0} = iz \frac{\partial}{\partial z} - i\bar{z} \frac{\partial}{\partial \bar{z}} + i \frac{\theta}{2} \frac{\partial}{\partial \theta} - i \frac{\bar{\theta}}{2} \frac{\partial}{\partial \bar{\theta}}, \quad (6.10)$$

$$X_{K_+} = iz^2 \frac{\partial}{\partial z} - i \frac{\partial}{\partial \bar{z}} + iz\theta \frac{\partial}{\partial \theta}, \quad X_{K_-} = i \frac{\partial}{\partial z} - i\bar{z}^2 \frac{\partial}{\partial \bar{z}} - i\bar{z}\bar{\theta} \frac{\partial}{\partial \bar{\theta}}, \quad (6.11)$$

$$X_{V_+} = z\theta \frac{\partial}{\partial z} + i \frac{\partial}{\partial \bar{\theta}}, \quad X_{W_-} = -i\bar{z}\bar{\theta} \frac{\partial}{\partial \bar{z}} - \frac{\partial}{\partial \theta}, \quad (6.12)$$

$$X_{V_-} = \theta \frac{\partial}{\partial z} + i\bar{z} \frac{\partial}{\partial \bar{\theta}}, \quad X_{W_+} = -i\bar{\theta} \frac{\partial}{\partial \bar{z}} - z \frac{\partial}{\partial \theta}. \quad (6.13)$$

This corresponds to simultaneously taking $b = -\tau$ and eliminating χ in (3.12).

From these explicit expressions one immediately identifies the superalgebra of the isotropy subsupergroup G_0 at the origin ($z = 0, \theta = 0$) of the atypical phase space. Clearly, X_B, X_{K_0}, X_{V_-} and X_{W_+} act trivially at the origin. Hence, the isotropy subsupergroup is generated by $\{B, K_0, V_-, W_+\}$. From the commutation relations (2.2) it appears that $G_0 \equiv \text{U}(1/1)$, and thus the atypical $\text{OSp}(2/2)$ -CS are parametrized by the super Kähler homogeneous space $\text{OSp}(2/2)/\text{U}(1/1) \equiv (\mathcal{D}^{(1|1)}, \omega)$. This could have been already deduced from (2.5)–(2.6) and (6.1).

Rothstein's data for $(\mathcal{D}^{(1|1)}, \omega)$ are given in [8]. They can be rederived by simply using results of Section 4. Indeed, the rank 2 holomorphic vector bundles $\mathbb{E}$ and $\mathbb{F}$ reduce both to a holomorphic line bundle. Moreover, from (4.7) one easily sees that when $b = -\tau$, we

are left with the Hermitian metric $||g|| = g_{1\bar{1}} = 2\tau/(1 - |z|^2)$. This is twice the metric obtained in [8].

On the other hand geometric quantization leads to the super Hilbert space obtained in [9]. This result corresponds to taking $b = -\tau$ in Section 5. Generators of $\mathfrak{osp}(2/2)$ are however represented by the following operators,

$$\widehat{B} = \frac{\theta}{2} \frac{\partial}{\partial \theta} - \tau, \quad \widehat{K}_0 = z \frac{\partial}{\partial z} + \frac{\theta}{2} \frac{\partial}{\partial \theta} + \tau, \quad (6.14)$$

$$\widehat{K}_+ = z^2 \frac{\partial}{\partial z} + z\theta \frac{\partial}{\partial \theta} + 2\tau z, \quad \widehat{K}_- = \frac{\partial}{\partial z}, \quad (6.15)$$

$$\widehat{V}_+ = -iz\theta \frac{\partial}{\partial z} - 2i\tau\theta, \quad \widehat{W}_- = i \frac{\partial}{\partial \theta}, \quad (6.16)$$

$$\widehat{V}_- = -i\theta \frac{\partial}{\partial z}, \quad \widehat{W}_+ = iz \frac{\partial}{\partial \theta}. \quad (6.17)$$

Finally, the supermeasure on $\mathcal{D}^{(1|1)}$ can not be obtained simply as the limit $b \rightarrow -\tau$ of (3.19). It has to be evaluated starting from (6.9) using the same technique as in Section 3.3. This leads to,

$$d\mu(z, \bar{z}, \theta, \bar{\theta}) = \frac{i}{\pi(1 - |z|^2)} \left[1 + i \frac{\bar{\theta}\theta}{1 - |z|^2} \right] dz d\bar{z} d\theta d\bar{\theta}. \quad (6.18)$$

7 Miscellaneous results and discussions

Here are gathered few consequences of the main results of the paper. Few other important points are discussed further.

7.1. Square integrability. When speaking about coherent states the first of their properties that comes to mind is the so-called resolution of the identity. In the non-super case this property reflects the square integrability of the unitary irreducible representation these special states belong to. Does this notion extend to the super case? The answer is yes. A simple computation based on (3.19), (2.27) and the Berezin integration leads to the following:

$$\int_{\mathcal{D}^{(1|2)}} |\bar{z}, \bar{\theta}, \bar{\chi}\rangle \langle \bar{z}, \bar{\theta}, \bar{\chi}| d\mu = \mathbb{I}. \quad (7.1)$$

Here $\mathbb{I} \equiv \mathbb{I}_{V(\tau, b)}$. A similar identity holds for the atypical CS. Hence, this allows a straightforward super extension of the definition of a square integrable representation. The above identity (7.1) provides a new argument that can be added to those already listed in [33] in order to justify calling the super unitary irreducible representations of $\mathcal{OSp}(2/2)$ considered here *discrete series representations*. As for usual CS, another immediate consequence

of (7.1) is that $\mathcal{H}_{\tau,b}$ is a reproducing super Hilbert space. This applies to both typical and atypical super Hilbert spaces.

7.2. Status of z . For simplicity the variable z was considered from the begining as a usual complex number (see (2.23)). The main reason behind this choice is to make the connection between the results of Sections 2 and 3, and Rothstein's approach to supersymplectic supergeometry, as described in Section 4, free of any change of coordinates. The same argument applies to the integrals over the unit disc $\mathcal{D}^{(1)}$ that appear in Section 5 (see (5.13)–(5.14)). Indeed, if the soul of z was different from zero, then those integrals and the complex geometry of Section 4 would be meaningless, unless a change of coordinates transforming z into a 'soulless' variable is performed. Hence, our initial choice prevents us from making any change of coordinates.

7.3. Whitney sum, Einstein-Kähler manifolds and Einstein-Hermitian vector bundles. The rank 2 holomorphic vector bundle $\mathbb{E}$ intervening in Rothstein's data for the typical orbits (see Theorem 4.3) is the Whitney sum of two holomorphic line bundles over the unit disc $\mathcal{D}^{(1)}$, i.e. $\mathbb{E} = \mathbb{E}_1 \oplus \mathbb{E}_2$. Independently, each component of $\mathbb{E}$ provides Rothstein's data for a super Kähler subsupermanifold of $(\mathcal{D}^{(1|2)}, \omega)$. More precisely, $(\mathcal{D}^{(1)}, \omega_0, \mathbb{E}_i, g_i, \nabla^{g_i})$, for $i = 1$ or 2 are such data for a $(1|1)$ -dimensional super Kähler supermanifold, denoted $\mathcal{D}_{\tau \pm b}^{(1|1)}$, where g_1 (resp. g_2) is the Hermitian metric on $\mathbb{E}_1$ (resp. $\mathbb{E}_2$) given in (4.7) as the first (resp. second) diagonal entry; and ∇^{g_i} are the associated Hermitian connections. These supermanifolds are $N = 1$ extensions of $(\mathcal{D}^{(1)}, \omega_0)$, the 2-superforms of which are obtained from (3.3)–(3.6) by setting $\chi = 0$ in the first instance and $\theta = 0$ in the second. These two super unit discs are not $\text{OSp}(2/2)$ -homogeneous spaces, unless $b = \pm\tau$. Indeed, when $b = \pm\tau$ they both become $\mathcal{D}^{(1|1)}$, the $N = 1$ super unit disc, which is a supersymplectic homogeneous space for both $\text{OSp}(2/2)$ and $\text{OSp}(1/2)$ as shown in Section 6. Since both $\mathbb{E}_1$ and $\mathbb{E}_2$ are trivial line bundles over $\mathcal{D}^{(1)}$, Rothstein's data for $\mathcal{D}_{\tau+b}^{(1|1)}$ and $\mathcal{D}_{\tau-b}^{(1|1)}$ differ only by the constant factor $\tau \pm b$ in front of the Hermitian structure on these bundles (see (4.7)).

The above observations lead to the following interesting picture. Let us denote by $\mathbf{R}_\lambda$ the data $(\mathcal{D}^{(1)}, \omega_0, \mathbb{E}_\lambda, g_\lambda, \nabla^{g_\lambda})$, where ω_0 is given in (3.4), $\mathbb{E}_\lambda$ a trivial holomorphic line bundle over $\mathcal{D}^{(1)}$, $\|g_\lambda\| = \lambda(1 - |z|^2)^{-1}$ (with $\lambda \geq 0$) a Hermitian structure on $\mathbb{E}_\lambda$, and ∇^{g_λ} the corresponding Hermitian connection. Then we have: **(i)** when λ is a multiple of τ , $\mathbf{R}_\lambda$ are just Rothstein's data for $\mathcal{D}^{(1|1)}$; **(ii)** the other $\text{OSp}(2/2)$ -coadjoint orbit, $(\mathcal{D}^{(1|2)}, \omega)$, admits as Rothstein's data $\mathbf{R}_{\lambda_1} \oplus \mathbf{R}_{\lambda_2}$ such that $\lambda_1 + \lambda_2 = 2\tau$, where the symbol ' $\oplus$ ' indicates that we have to take the Whitney sum of $(\mathbb{E}_{\lambda_1}, g_{\lambda_1})$ and $(\mathbb{E}_{\lambda_2}, g_{\lambda_2})$. We recall

at this point that the information in $\mathbf{R}_{\tau-b} \oplus \mathbf{R}_{\tau+b}$ are sufficient to reconstruct the 2-superform (3.3)–(3.6) (see Section 4). Hence, $\mathbf{R}_\lambda$ is the basic building block for describing super Kähler coadjoint orbits of simple Lie supergroups super extending $SU(1, 1)$.

A deeper analysis of $\mathbf{R}_\lambda = (\mathcal{D}^{(1)}, \omega_0, \mathbb{E}_\lambda, g_\lambda, \nabla^{g_\lambda})$ shows that it is made of two main parts: **(i)** $(\mathcal{D}^{(1)}, \omega_0)$, which is a Kähler $SU(1, 1)$ -homogeneous space, and **(ii)** $(\mathbb{E}_\lambda, g_\lambda)$, which is an *Einstein-Kähler manifold* [45]. Here we identify $\mathbb{E}_\lambda$ with the holomorphic tangent bundle over $\mathcal{D}^{(1)}$. On the other hand, it appears that $\mathbf{R}_{\lambda_1} \oplus \mathbf{R}_{\lambda_2}$ defines an *Einstein-Hermitian vector bundle* [45]. These very interesting and important observations not only improve the characterization of a super Kähler coadjoint orbit, but they suggest a way of extending to the super context the known classification of irreducible bounded symmetric Hermitian domains. This direction is now under investigation.

7.4. Realizations of the typical and atypical representations. In agreement with the descriptions of Section 2 and Appendix B, results of Section 5 show that the typical super Hilbert space $\mathcal{H}_{\tau,b}$ of L^2 superholomorphic sections of $\mathcal{L}^{(1|2)}$ is the direct sum of Hilbert spaces of four holomorphic discrete series representations of $SU(1, 1)$. More precisely, as a vector superspace

$$\mathcal{H}_{\tau,b} = \mathcal{H}_{k=\tau} \oplus 2 \cdot \mathcal{H}_{k=\tau+\frac{1}{2}} \oplus \mathcal{H}_{k=\tau+1}, \quad (7.2)$$

where $\mathcal{H}_k$ is the Hilbert space carrying the holomorphic discrete series representation $D(k)$ of $SU(1, 1)$ (see (5.14) and (2.8)–(2.10)). This suggests that the $\mathfrak{osp}(2/2)$ operators obtained in (5.17) and which act in the left hand side of (7.2) can be replaced by matrix valued and thus anticommutating-variables free operators acting in the right hand side of (7.2) [55]. The former realization is much more convenient than the latter. Indeed, for example for $\mathfrak{osp}(N/2)$, the matrices can be $2^N \times 2^N$. We insist here on the fact that our main goal in Section 5 was to show that geometric quantization extends to the super context, at least when applied to coadjoint orbits admitting a super Kähler polarization. We not only succeeded in achieving this, but the above observation confirms that our output constitutes an intrinsically supersymmetric alternative to the matrix realization. The same discussion applies to the atypical representations.

7.5. $OSp(2/2)$ representations. Throughout, we have been considering only representations of the Lie superalgebra. Explicit representations of $OSp(2/2)$ can be in fact obtained from those of $\mathfrak{osp}(2/2)$ exhibited in Sections 5 and 6. The procedure does not present any difficulties. The first step towards this construction consists in finding the explicit action of $OSp(2/2)$ on $\mathcal{D}^{(1|2)}$ and $\mathcal{D}^{(1|1)}$, by integrating the Hamiltonian vector fields in (3.12)

and (6.10)–(6.13). This amounts to solving super Riccati differential equations, which have already been considered in [56]. The full construction will be given elsewhere.

8 Conclusions and outlook

Although our present contribution treats a specific example, the obtained results pave the way to harmonic superanalysis. It must be regarded as the first important step of a program aimed at classifying Lie supergroups' coadjoint orbits and the associated irreducible representations.

In this work several closely related questions have been addressed, and several new notions have been introduced. The consistency of our conventions is manifest throughout the paper, from abstract to explicit representation theory via super Kähler geometry. The main results are now summarized:

- (a) Starting with a comprehensive description of the abstract typical and atypical representations of $\mathfrak{osp}(2/2)$, the associated $\mathrm{OSp}(2/2)$ coherent states are constructed.
- (b) Super extending known methods, their underlying geometries are exhibited, and shown to be those of $\mathrm{OSp}(2/2)$ coadjoint orbits. The latter are $\mathrm{OSp}(2/2)$ -supersymplectic homogeneous spaces: $\mathcal{D}^{(1|2)} \equiv \mathrm{OSp}(2/2)/(\mathrm{U}(1) \times \mathrm{U}(1))$ for the typical CS, and $\mathcal{D}^{(1|1)} \equiv \mathrm{OSp}(2/2)/\mathrm{U}(1/1)$ for the atypical CS.
- (c) The identification of Rothstein's data for $\mathcal{D}^{(1|2)}$ and $\mathcal{D}^{(1|1)}$, draws us to generalizing Rothstein's theorem to the complex-analytic setting. This leads to a natural definition of a super Kähler supermanifold, $\mathcal{D}^{(1|2)}$ and $\mathcal{D}^{(1|1)}$ being non-trivial examples of such a notion. We moreover show that in this context, Rothstein's theorem can be refined. More precisely, the *complete* supersymplectic structure of a super Kähler coadjoint orbit can be encoded in an elementary building block of the type mentioned in point 7.3 of the previous section.
- (d) Finally, geometric quantization is successfully extended to the super Kähler context exemplified by the typical and atypical coadjoint orbits of $\mathrm{OSp}(2/2)$. A super Kähler polarization is exhibited in each case. This leads to an explicit super unitary irreducible typical (atypical) representation of $\mathfrak{osp}(2/2)$ in a super Hilbert space of square integrable superholomorphic sections of a complex line bundle sheaf over $\mathcal{D}^{(1|2)}$ ($\mathcal{D}^{(1|1)}$).

Possible generalizations of our results are numerous and worth considering. At both the representation theoretic and the geometric levels, the present work relies essentially on known results from the non-super context. For instance, the representation theory of

$\text{osp}(2/2)$ is based on that of $\text{su}(1,1) \subset \text{osp}(2/2)$, while the supergeometry of the $N = 2$ (resp. $N = 1$) super unit disc $\mathcal{D}^{(1|2)}$ (resp. $\mathcal{D}^{(1|1)}$) is based on that of the unit disc $\mathcal{D}^{(1)}$. Our results show precisely how the super extension occurs (see Section 7, 7.3). One can now seriously consider other Lie supergroups, and look for a classification of their super Kähler homogeneous spaces along the known classification of the Kähler homogeneous spaces of their body Lie groups. Geometric quantization will then provide a classification of their associated representations.

Acknowledgments A.M.E. is indebted to S.T. Ali, C. Duval, J. Harnad, and G. Tuynman for valuable and stimulating discussions. He thanks the Mathematics Department of Concordia University, the Centre de Recherches Mathématiques of Université de Montréal, and the U.F.R. de Mathématiques of Université de Lille for their kind hospitality, and much appreciated support. L.M.N. acknowledges a fellowship from Ministerio de Educación y Ciencia (Spain) and kind hospitality at the Centre de Recherches Mathématiques of Université de Montréal.

A Conventions and notation

One of the main features of the present work is the complete consistency of the conventions used throughout. The latter are displayed here. They concern the superalgebra $\mathcal{B}$, introduced in Section 2, and its interactions with both the Lie superalgebra and the Lie superalgebra's modules.

A complex superalgebra is a complex vector superspace (i.e. a $\mathbb{Z}_2$ -graded linear space) $\mathcal{B} = \mathcal{B}_0 \oplus \mathcal{B}_1$ equipped with a $\mathbb{Z}_2$ -compatible product, namely, $\mathcal{B}_k \cdot \mathcal{B}_l \subset \mathcal{B}_{k+l}$; $\mathcal{B}$ is considered associative and possesses a unity. Note that, $\mathcal{B}_0$ (resp. $\mathcal{B}_1$) is called the *even* (resp. *odd*) part of $\mathcal{B}$. Accordingly, elements of $\mathcal{B}_0$ (resp. $\mathcal{B}_1$) are called *even* (resp. *odd*) elements of $\mathcal{B}$. A homogeneous element of $\mathcal{B}$ is either even or odd. The *parity* of such an element $u \in \mathcal{B}_k$, denoted $\epsilon(u)$, is defined by $\epsilon(u) = k$. The superalgebra $\mathcal{B}$ is supercommutative if

$$uv = (-1)^{\epsilon(u)\epsilon(v)}vu, \quad (\text{A.1})$$

for u and v two homogeneous elements of $\mathcal{B}$.

The *complex supercommutative superalgebra with unit* $\mathcal{B}$ considered in the present work is the complex Grassmann algebra [22, 23] generated by (θ, χ) and their complex conjugates $(\bar{\theta}, \bar{\chi})$. These are anticommuting, and hence nilpotent variables. In other

words $\mathcal{B}$ is the complex exterior algebra over $\mathbb{C}^4 = \mathbb{C}^2 \oplus \overline{\mathbb{C}}^2$. Its even (resp. odd) part is spanned by the products of an even (resp. odd) number of generators, and the dimension of $\mathcal{B}$ is 16. The decomposition of any element $\Theta \in \mathcal{B}$ in a given basis of $\mathcal{B}$, assumes the following form

$$\Theta = \tilde{\Theta} \cdot \mathbf{I} + \Theta_{nil}, \quad (\text{A.2})$$

where, the purely nilpotent component Θ_{nil} is called the *soul* of Θ , while the component $\tilde{\Theta}$ along the identity of $\mathcal{B}$ is called the *body* of Θ .

The complex conjugation maps $\mathbb{C}^2 \ni (\theta, \chi) \mapsto (\bar{\theta}, \bar{\chi}) \in \overline{\mathbb{C}}^2$. Its extension to $\mathcal{B}$ is completely defined by the following rule:

$$\overline{\Theta_1 \Theta_2} = \bar{\Theta}_1 \bar{\Theta}_2, \quad \forall \Theta_1, \Theta_2 \in \mathcal{B}. \quad (\text{A.3})$$

The other properties are:

$$\bar{\bar{\Theta}} = \Theta \quad \text{and} \quad \overline{w\Theta} = \bar{w}\bar{\Theta}, \quad \forall w \in \mathbb{C} \text{ and } \Theta \in \mathcal{B}. \quad (\text{A.4})$$

An element $\Theta \in \mathcal{B}$ is *real* if $\bar{\Theta} = \Theta$. Using (A.3) one easily sees that $\Theta\bar{\Theta}$ is real for $\Theta \in \mathcal{B}_0$ and imaginary for $\Theta \in \mathcal{B}_1$.

It is important to notice that our convention in (A.3) is different from the one introduced by Berezin [21, 22], and commonly used in the literature (see [53] and references therein). In that case the complex conjugate of a product is taken as:

$$\overline{\Theta_1 \Theta_2} = \bar{\Theta}_2 \bar{\Theta}_1 \quad \forall \Theta_1, \Theta_2 \in \mathcal{B}. \quad (\text{A.5})$$

Hence, for Θ a homogeneous element of $\mathcal{B}$, $\Theta\bar{\Theta}$ is real independently of the parity of Θ . Using these conventions one faces serious inconsistencies. The most obvious one was encountered in [8] (see also [50]), where the author followed Berezin's conventions already used in [34]; the super Kähler 2-form obtained there was neither real nor imaginary! As a consequence, the classical observables and their associated Hamiltonian vector fields were not satisfying any property of the type of (3.10) and (3.13), which are crucial in identifying a real observable and then the associated self superadjoint operator. When combined with the notion of a super Hermitian structure of [41], our convention (A.3) cures this discrepancy. More precisely, the arguments invoked in [53] in order to justify the choice in (A.5) apply to our choice (A.3) too, provided one considers the notion of a super Hermitian structure (2.12)–(2.16) and its consequences. (The problem in [8] mentioned above is cured in [9]).

Finally, all vector superspaces appearing in this work are considered as left $\mathcal{B}$ -modules. Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ be such a $\mathcal{B}$ -module. Then, for v and Θ homogeneous elements in V and $\mathcal{B}$ respectively, we have,

$$\Theta v = (-1)^{\epsilon(\Theta)\epsilon(v)} v \Theta. \quad (\text{A.6})$$

This applies equally well to $V = \text{osp}(2/2, \mathbb{C})$, $V = V(\tau, b)$ or $V = U(\pm\tau)$. When V is equipped with an additional structure, such as a super Hermitian form $\langle \cdot | \cdot \rangle$ (see (2.12)–(2.14)) or a Lie superalgebra bracket $[\cdot, \cdot]$, then that structure can be extended to the *Grassmann envelope of second type* $\tilde{V}$ of V . The latter is defined as follows [22],

$$\tilde{V} \equiv (\mathcal{B} \otimes V)_0 = (\mathcal{B}_0 \otimes V_0) \oplus (\mathcal{B}_1 \otimes V_1), \quad (\text{A.7})$$

and the structures above are extended in the following way:

$$\langle \Theta_1 v | \Theta_2 u \rangle = (-1)^{\epsilon(v)\epsilon(\Theta_2)} \bar{\Theta}_1 \Theta_2 \langle v | u \rangle \quad \text{and} \quad [\Theta_1 X, \Theta_2 Y] = (-1)^{\epsilon(X)\epsilon(\Theta_2)} \Theta_1 \Theta_2 [X, Y], \quad (\text{A.8})$$

where Θ_2 , v and X are homogeneous elements in respectively, $\mathcal{B}$, $V(\tau, b)$ and $\text{osp}(2/2, \mathbb{C})$. We end this appendix by giving a formula which is useful for some of the computations of Section 2 and Appendix B. Let Θ and v be homogeneous elements of respectively $V(\tau, b)$ and $\mathcal{B}$, the super Hermitian conjugate of $|\Theta v\rangle \in \tilde{V}(\tau, b)$ with respect to the super Hermitian form (2.13)–(2.16), is obtained as follows:

$$(\Theta | v \rangle)^\dagger = \bar{\Theta} (| v \rangle)^\dagger = (i)^{\epsilon(v)} \bar{\Theta} \langle v|. \quad (\text{A.9})$$

B $\text{osp}(2/2)$ representations: more details

Finite and infinite dimensional irreducible representations of $\text{osp}(2/2)$ have already been studied in [32, 33, 35]. The description of the infinite dimensional ones given in [35] is the most convenient for our purpose, but since it suffers from some discrepancies we consider important to reexpose the construction. This appendix must be viewed as a complement to Section 2.

The equations defining the lowest weight vector $|0\rangle$ (2.5)–(2.6), together with the observation that $|0\rangle$ is the lowest weight vector of a discrete series representation $D(\tau)$ of $\text{su}(1, 1)$ (2.7), are our starting points.

The $\text{osp}(2/2)$ -module $V(\tau, b)$ of Section 2 is generated by applying arbitrary polynomials in the generators of $\mathfrak{n}^+ = \text{span}\{K_+, V_+, W_+\}$ (2.3) to $|0\rangle$. Using the commutation

relations and the results mentioned above, one can see that $V(\tau, b)$ is spanned by the following vectors:

$$K_+^m|0\rangle, \quad K_+^m V_+|0\rangle, \quad K_+^m W_+|0\rangle, \quad K_+^m V_+ W_+|0\rangle, \quad m \in \mathbb{N}. \quad (\text{B.1})$$

The latter are eigenstates of B :

$$\begin{aligned} B(K_+^m|0\rangle) &= b(K_+^m|0\rangle) \\ B(K_+^m V_+|0\rangle) &= (b + \tfrac{1}{2})(K_+^m V_+|0\rangle) \\ B(K_+^m W_+|0\rangle) &= (b - \tfrac{1}{2})(K_+^m W_+|0\rangle) \\ B(K_+^m V_+ W_+|0\rangle) &= b(K_+^m V_+ W_+|0\rangle), \end{aligned} \quad (\text{B.2})$$

and also of K_0 :

$$\begin{aligned} K_0(K_+^m|0\rangle) &= (\tau + m)(K_+^m|0\rangle) \\ K_0(K_+^m V_+|0\rangle) &= (\tau + \tfrac{1}{2} + m)(K_+^m V_+|0\rangle) \\ K_0(K_+^m W_+|0\rangle) &= (\tau + \tfrac{1}{2} + m)(K_+^m W_+|0\rangle) \\ K_0(K_+^m V_+ W_+|0\rangle) &= (\tau + 1 + m)(K_+^m V_+ W_+|0\rangle), \end{aligned} \quad (\text{B.3})$$

but not all of them are eigenstates of the $\text{su}(1, 1)$ Casimir C_2 given in (2.18):

$$\begin{aligned} C_2(K_+^m|0\rangle) &= \tau(\tau - 1)(K_+^m|0\rangle) \\ C_2(K_+^m V_+|0\rangle) &= (\tau + \tfrac{1}{2})(\tau - \tfrac{1}{2})(K_+^m V_+|0\rangle) \\ C_2(K_+^m W_+|0\rangle) &= (\tau + \tfrac{1}{2})(\tau - \tfrac{1}{2})(K_+^m W_+|0\rangle) \\ C_2(K_+^m V_+ W_+|0\rangle) &= (\tau + 1)\tau(K_+^m V_+ W_+|0\rangle) - (\tau + b)(K_+^{m+1}|0\rangle). \end{aligned} \quad (\text{B.4})$$

The last equation suggests to use another family of states instead of $K_+^m V_+ W_+|0\rangle$. Indeed, notice that the vectors

$$(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle, \quad (\text{or} \quad (-2\tau K_+^m W_+ V_+ + (\tau - b)K_+^{m+1})|0\rangle), \quad (\text{B.5})$$

are such that

$$\begin{aligned} B(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle &= b(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle \\ C_2(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle &= (\tau + 1)\tau(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle \\ K_0(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle &= (\tau + 1 + m)(2\tau K_+^m V_+ W_+ - (\tau + b)K_+^{m+1})|0\rangle. \end{aligned} \quad (\text{B.6})$$

The previous results suggest to use the following notation:

$$\begin{aligned} |b, \tau, \tau\rangle &\equiv |0\rangle, & |b + \tfrac{1}{2}, \tau + \tfrac{1}{2}, \tau + \tfrac{1}{2}\rangle &\propto V_+|0\rangle, \\ |b, \tau + 1, \tau + 1\rangle &\propto (2\tau V_+ W_+ - (\tau + b)K_+)|0\rangle, & |b - \tfrac{1}{2}, \tau + \tfrac{1}{2}, \tau + \tfrac{1}{2}\rangle &\propto W_+|0\rangle. \end{aligned} \quad (\text{B.7})$$

The $\text{osp}(2/2)$ -module $V(\tau, b)$ obtained in this way is irreducible only when $b \neq \pm\tau$. For $b = \pm\tau$, $V(\tau, b)$ contains a *primitive vector*. Indeed, using (2.2a)–(2.2i) and (2.5)–(2.6), one easily sees that:

$$\text{when } b = \tau, \quad K_-(V_+|0\rangle) = V_-(V_+|0\rangle) = W_-(V_+|0\rangle) = 0, \quad (\text{B.8})$$

and

$$\text{when } b = -\tau, \quad K_-(W_+|0\rangle) = V_-(W_+|0\rangle) = W_-(W_+|0\rangle) = 0. \quad (\text{B.9})$$

These two situations being very similar, we focus here only on the second one. Hence, in that instance, $W_+|0\rangle$ generates an $\text{osp}(2/2)$ -submodule of $V(\tau, b)$, denoted $V'(\tau, -\tau)$. An irreducible $\text{osp}(2/2)$ -module emerges then as the quotient $V(\tau, -\tau)/V'(\tau, -\tau) \equiv U(-\tau)$. Notice that $V'(\tau, -\tau)$ is spanned by the last two series in (B.1), while $U(-\tau)$ is spanned by the two first ones modulo the two last ones.

In order to obtain the proportionality constants in (B.7) it is necessary to equip $V(\tau, b)$ with a super Hermitian form [41, 24] of the type described in (2.12)–(2.14). The *superadjoint* $A^\dagger$ (denoted $\tilde{A}$ and called differently in [24]) of a homogeneous operator A acting in $V(\tau, b)$ is defined as follows:

$$\langle A^\dagger u | v \rangle = (-1)^{\epsilon(u)\epsilon(A)} \langle u | Av \rangle, \quad \forall u, v \in V(\tau, b) \quad \text{with } u \text{ homogeneous.} \quad (\text{B.10})$$

Hence, A is self superadjoint if $A^\dagger = A$. Moreover, one can check from (B.10) that

$$(A^\dagger)^\dagger = A \quad \text{and} \quad (AB)^\dagger = (-1)^{\epsilon(A)\epsilon(B)} B^\dagger A^\dagger. \quad (\text{B.11})$$

Consequently,

$$[A, B]^\dagger = -[A^\dagger, B^\dagger]. \quad (\text{B.12})$$

Using this equation, the relations in (2.2a)–(2.2i) and the root space decomposition of $\text{osp}(2/2, \mathbb{C})$ given in Section 2, one easily shows that

$$B^\dagger = B, \quad K_0^\dagger = K_0, \quad (K_\pm)^\dagger = K_\mp, \quad (\text{B.13})$$

$$(V_\pm)^\dagger = iW_\mp \quad \text{and} \quad (W_\pm)^\dagger = iV_\mp. \quad (\text{B.14})$$

Using the above results, equation (A.9), and assuming that the vectors on the left hand side of the equations in (B.7) are normalized to one with respect to $\langle \cdot | \cdot \rangle_{0(1)}$ (according to their parity) one obtains:

$$V_+|b, \tau, \tau\rangle = \sqrt{\tau - b}|b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2}\rangle, \quad (\text{B.15})$$

$$W_+|b, \tau, \tau\rangle = \sqrt{\tau + b}|b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2}\rangle, \quad (\text{B.16})$$

$$(2\tau V_+ W_+ - (\tau + b)K_+) |b, \tau, \tau\rangle = \sqrt{(\tau^2 - b^2)(2\tau + 1)2\tau} |b, \tau + 1, \tau + 1\rangle. \quad (\text{B.17})$$

Clearly, these results are valid only if $|b| \leq \tau$. When $b \neq \pm\tau$, this means that the super Hermitian structure introduced above, turns the irreducible $\text{osp}(2/2)$ -modules $V(\tau, b)$ into super unitary representations only if $|b| < \tau$. On the other hand, when $b = \pm\tau$ one sees from (B.15)–(B.17) that the primitive vectors generating the submodules $V'(\tau, \pm\tau)$ are zero-norm states, and thus the entire submodules are made of zero-norm states. Moding out the latter from $V(\tau, \pm\tau)$ turns $U(\pm\tau)$ into super unitary irreducible modules. Moreover, (2.9) and (2.11) are direct consequences of (B.15)–(B.17) and (2.2a)–(2.2i). Indeed, the four states $|b, \tau, \tau\rangle$, $V_+|b, \tau, \tau\rangle$, $W_+|b, \tau, \tau\rangle$ and $(2\tau V_+ W_+ - (\tau + b)K_+) |b, \tau, \tau\rangle$ are orthogonal with respect to the super Hermitian structure, and each of them is a lowest state of an $\text{su}(1, 1)$ irreducible module. The latter are generated from the former $\text{su}(1, 1)$ lowest-weight states through the action of powers of K_+ :

$$K_+^m | \cdot, k, k \rangle = \sqrt{\frac{m! \Gamma(2k + m)}{\Gamma(2k)}} | \cdot, k, k + m \rangle; \quad k = \tau, \tau \pm \frac{1}{2}, \text{ or } \tau + 1. \quad (\text{B.18})$$

We end this appendix by displaying the action of the $\text{osp}(2/2)$ generators on the different vectors. The following formulae are straightforward consequences of (2.2a)–(2.2i) and the results described above;

$$\begin{aligned} K_+|b, \tau, \tau + m\rangle &= \sqrt{(2\tau + m)(m + 1)} |b, \tau, \tau + m + 1\rangle, \\ K_-|b, \tau, \tau + m\rangle &= \sqrt{(2\tau + m - 1)m} |b, \tau, \tau + m - 1\rangle, \\ V_+|b, \tau, \tau + m\rangle &= \sqrt{\frac{(\tau - b)(2\tau + m)}{2\tau}} |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle, \\ V_-|b, \tau, \tau + m\rangle &= \sqrt{\frac{(\tau - b)m}{2\tau}} |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m - 1\rangle, \\ W_+|b, \tau, \tau + m\rangle &= \sqrt{\frac{(\tau + b)(2\tau + m)}{2\tau}} |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle, \\ W_-|b, \tau, \tau + m\rangle &= \sqrt{\frac{(\tau + b)m}{2\tau}} |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m - 1\rangle, \\ V_+|b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= 0, \end{aligned}$$

$$\begin{aligned}
V_-|b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= 0, \\
W_+|b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= \sqrt{\frac{(\tau-b)(m+1)}{2\tau}} |b, \tau, \tau + m + 1\rangle \\
&\quad - \sqrt{\frac{(\tau+b)(2\tau+m+1)}{2\tau}} |b, \tau + 1, \tau + 1 + m\rangle, \\
W_-|b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= \sqrt{\frac{(\tau-b)(2\tau+m)}{2\tau}} |b, \tau, \tau + m\rangle \\
&\quad - \sqrt{\frac{(\tau+b)m}{2\tau}} |b, \tau + 1, \tau + 1 + m - 1\rangle, \\
V_+|b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= \sqrt{\frac{(\tau+b)(m+1)}{2\tau}} |b, \tau, \tau + m + 1\rangle \\
&\quad + \sqrt{\frac{(\tau-b)(2\tau+m+1)}{2\tau}} |b, \tau + 1, \tau + 1 + m\rangle, \\
V_-|b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= \sqrt{\frac{(\tau+b)(2\tau+m)}{2\tau}} |b, \tau, \tau + m\rangle \\
&\quad + \sqrt{\frac{(\tau-b)m}{2\tau}} |b, \tau + 1, \tau + 1 + m - 1\rangle, \\
W_+|b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= 0, \\
W_-|b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle &= 0, \\
V_+|b, \tau + 1, \tau + 1 + m\rangle &= -\sqrt{\frac{(\tau+b)(m+1)}{2\tau}} |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m + 1\rangle, \\
V_-|b, \tau + 1, \tau + 1 + m\rangle &= -\sqrt{\frac{(\tau+b)(2\tau+m+1)}{2\tau}} |b + \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle, \\
W_+|b, \tau + 1, \tau + 1 + m\rangle &= \sqrt{\frac{(\tau-b)(m+1)}{2\tau}} |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m + 1\rangle, \\
W_-|b, \tau + 1, \tau + 1 + m\rangle &= \sqrt{\frac{(\tau-b)(2\tau+m+1)}{2\tau}} |b - \frac{1}{2}, \tau + \frac{1}{2}, \tau + \frac{1}{2} + m\rangle.
\end{aligned} \tag{B.19}$$

The discrepancies mentioned at the beginning of this appendix can be easily seen by comparing (B.19) with its analog in [35].

References

1. E. Schrödinger, *Der stetige Übergang von der Mikro- zur Makromechanik*, Naturwissenschaften **14** (1926) 664.
2. R.J. Glauber, *The quantum theory of optical coherence*, Phys. Rev. **130** (1963) 2529; *Coherent and incoherent states of the radiation field*, Phys. Rev. **131** (1963) 2766.
3. A. Perelomov, *Generalized Coherent States and Their Applications* (Springer-Verlag, Berlin, 1986).
4. J.R. Klauder and B.-S. Skagerstam, *Coherent States – Applications in Physics and Mathematical Physics* (World Scientific, Singapore, 1985).
5. I. Daubechies, *Ten Lectures on Wavelets*, CBMS-NSF Regional Conference Series in Applied Mathematics **61** (SIAM, Philadelphia, 1992).
6. F.A. Berezin, *General concept of quantization*, Commun. Math. Phys. **40** (1975) 153.

7. E. Onofri, *A note on coherent state representations of Lie groups*, J. Math. Phys. **16** (1975) 1087.
8. A.M. El Gradechi, *On the supersymplectic homogeneous superspace underlying the $OSp(1/2)$ coherent states*, J. Math. Phys. **34** (1993) 5951.
9. A.M. El Gradechi, *Supergeometric quantization of an $OSp(1/2)$ coadjoint orbit*. Preprint CRM-1882. Submitted.
10. W.M. Zhang, D.H. Feng, and R. Gilmore, *Coherent states: Theory and some applications*, Rev. Mod. Phys. **26** (1990) 867.
11. R. Delbourgo, *Minimal uncertainty states for the rotation and allied groups*, J. Phys. A: Math. Gen. **10** (1977) 1837; R. Delbourgo and J. R. Fox, *Maximum weight vectors possess minimal uncertainty*, J. Phys. A: Math. Gen. **10** (1977) L235.
12. S. De Bièvre and A.M. El Gradechi, *Quantum mechanics and coherent states on the anti-de Sitter spacetime and their Poincaré contraction*, Ann. Inst. H. Poincaré **57** (1992) 403.
13. J.M. Radcliffe, *Some properties of spin coherent states*, J. Phys. A: Gen. Phys. **4** (1971) 313.
14. S.T. Ali and J.-P. Antoine, *Coherent states of the 1 + 1-dimensional Poincaré group: square integrability and a relativistic Weyl transform*, Ann. Inst. H. Poincaré **51** (1989) 23; S.T. Ali, J.-P. Antoine and J.-P. Gazeau, *De Sitter to Poincaré contraction and relativistic coherent states*, Ann. Inst. H. Poincaré **52** (1990) 83.
15. S.T. Ali, J.-P. Antoine and J.-P. Gazeau, *Square integrability of group representations on homogeneous spaces. I. Reproducing triples and frames*, Ann. Inst. H. Poincaré **55** (1991) 829; *II. Coherent and quasi-coherent states. The case of the Poincaré group*, Ann. Inst. H. Poincaré **55** (1991) 857.
16. A. Kirillov, *Eléments de la Théorie des Représentations*, (Editions Mir, Moscou 1974); *The method of orbits in representation theory*, in: Lie Groups and Their Representations, I.M. Gelfand (Ed.), (Akadémiai Kiadó, Budapest 1975), p. 219.
17. B. Kostant, *Quantization and unitary representations*, in: Lecture Notes in Mathematics **170** (Springer-Verlag, New York 1970), p. 87.
18. J.M. Souriau, *Structure des Systèmes Dynamique* (Dunod, Paris, 1970).
19. F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, and D. Sternheimer, *Deformation theory and quantization*, Ann. Phys. (N.Y.) **111** (1978) 61.
20. P.G.O. Freund, *Introduction to Supersymmetry* (Cambridge University Press, Cambridge, 1986).
21. F.A. Berezin, *The Method of Second Quantization* (Academic Press, New York, 1966).
22. F.A. Berezin, *Introduction to Superanalysis* (Reidel, Dordrecht, 1987); *The mathematical basis of supersymmetric field theories*, Soviet J. Nucl. Phys. **29** (1979) 857.
23. J.F. Cornwell, *Group Theory in Physics Vol.3 – Supersymmetries and Infinite Dimensional Algebras* (Academic Press, London, 1989).
24. M. Scheunert, *The Theory of Lie Superalgebras – An Introduction*, Lecture Notes in Mathematics **716** (Springer-Verlag, Berlin, 1979).
25. V. Kac, *Representations of classical Lie superalgebras* in: Lecture Notes in Mathematics **676** (Springer-Verlag, Berlin, 1978) p.597.
26. D.A. Leites, *Introduction to the theory of supermanifolds*, Russian Math. Surveys **35** (1980) 1.
27. B.S. DeWitt, *Supermanifolds* 2nd Edition (Cambridge University Press, Cambridge, 1992).

28. B. Kostant, *Graded manifolds, graded Lie theory and prequantization*, in: Conference on Differential Geometric Methods in Mathematical Physics (K. Bleuler and A. Reetz eds.), Lecture Notes in Mathematics **570** (Springer-Verlag, Berlin, 1977) p. 177.
29. Y.I. Manin, *Gauge Field Theory and Complex Geometry* (Springer-Verlag, Berlin, 1988).
30. C. Bartocci, U. Bruzzo and D. Hernández-Ruipérez, *The geometry of Supermanifolds* (Kluwer Academic Publishers, Dordrecht, 1991).
31. G.M. Tuynman, *Supermanifolds: A Geometric Approach*. Book in preparation.
32. M. Scheunert, W. Nahm, V. Rittenberg, *Irreducible representations of the $osp(2,1)$ and $spl(2,1)$ graded Lie algebras*, J. Math. Phys. **18** (1977) 155.
33. K. Nishiyama, *Characters and super-characters of discrete series representations for orthosymplectic Lie superalgebras*, J. Algebra **141** (1991) 399.
34. A.B. Balantekin, H.A. Schmitt and B.R. Barrett, *Coherent states for the harmonic oscillator representations of the orthosymplectic supergroup $OSp(1/2N, R)$* , J. Math. Phys. **29** (1988) 1634.
35. A.B. Balantekin, H.A. Schmitt and P. Halse, *Coherent states for the noncompact supergroups $OSp(2/2N, R)$* , J. Math. Phys. **30** (1989) 274.
36. B.W. Fatyga, V.A. Kostelecký, M.M. Nieto and D.R. Truax, *Supercoherent states*, Phys. Rev. **D43** (1991) 1403.
37. C. Duval and P.A. Horváthy, *On Schrödinger superalgebras*, Preprint CPT-93/P.2912 (1993).
38. M. Rothstein, *The structure of supersymplectic supermanifolds*, in: Differential Geometric Methods in Mathematical Physics (C. Bartocci, U. Bruzzo, and R. Cianci eds.), Lecture Notes in Physics **375** (Springer-Verlag, Berlin, 1991) p. 331.
39. J. Monterde, *A characterization of graded symplectic structures*, Differential Geom. Appl. **2** (1992) 81.
40. L. Frappat, A. Sciarrino, and P. Sorba, *Structure of basic Lie superalgebras and of their affine extensions*, Commun. Math. Phys. **121** (1989) 457, and references therein.
41. S. Sternberg and J. Wolf, *Hermitian Lie algebras and metaplectic representations*, Trans. Amer. Math. Soc. **238** (1978) 1.
42. P. Green, *On holomorphic graded manifolds*, Proc. Amer. Math. Soc. **85** (1982) 587.
43. M. Rothstein, *Deformations of complex supermanifolds*, Proc. Amer. Math. Soc. **95** (1985) 255.
44. M. Batchelor, *The structure of supermanifolds*, Trans. Amer. Math. Soc. **253** (1979) 329.
45. S. Kobayashi, *Differential Geometry of Complex Vector Bundles* (Princeton University Press, Princeton, 1987).
46. N.M.J. Woodhouse, *Geometric Quantization* (Clarendon Press, Oxford, 1980).
47. A.A. Kirillov, *Geometric quantization*, in: Dynamical Systems IV, (V.I. Arnol'd and S.P. Novikov eds., Springer-Verlag, Berlin, 1990) p. 137.
48. G.M. Tuynman, *Geometric quantization of the BRST charge*, Commun. Math. Phys. **150** (1992) 237.
49. P. Schaller and G. Schwarz, *Anomalies from geometric quantization of fermionic field theories*, J. Math. Phys. **31** (1990) 2366.
50. D. Borthwick, S. Klimek, A. Lesniewski, and M. Rinaldi, *Super Toeplitz operators and non-*

- perturbative quantization of supermanifolds*, Commun. Math. Phys. **153** (1993) 49.
51. G.M. Tuynman, *Quantization: Towards a comparison between methods*, J. Math. Phys. **28** (1987) 2829.
 52. P. Renouard, *Variétés Symplectiques et Quantification*, Thesis, Orsay, 1969.
 53. T. Schmitt, *Supergeometry and hermitian conjugation*, J. Geom. Phys. **7** (1990) 141.
 54. S. Nagamachi and Y. Kobayashi, *Hilbert superspace*, J. Math. Phys. **33** (1992) 4274.
 55. A.M. El Gradechi, *Matrix-valued operators versus superoperators: the orthosymplectic Lie superalgebras*, in preparation.
 56. J. Beckers, L. Gagnon, V. Hussin, and P. Winternitz, *Superposition formulas for nonlinear super-equations*, J. Math. Phys. **31** (1990) 2528.